

Please type a sign(+) inside this box



PTO/SB/05 (2/98)

Approved for use through 09/30/2000 OMB 0651-0032
Patent and Trademark Office U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR § 1.53(b))

Attorney Docket No.	0942.4330003/RWE/BJD
First Inventor or Application Identifier	GERARD
Title	Compositions and Methods for Reverse Transcription of Nucleic Acid Molecules
Express Mail Label No.	

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1. ☒ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 133]
(preferred arrangement set forth below)
 - Descriptive title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
3. ☒ Drawing(s) (35 U.S.C. 113) [Total Sheets 54]
4. ☒ Oath or Declaration [Total Pages 2]
 - a. ☐ Newly executed (original or copy)
 - b. ☒ Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional with Box 17 completed) [Note Box 5 below]
 - i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR §§ 1.63(d)(2) and 1.33(b).
5. ☒ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein
6. ☐ Microfiche Computer Program (Appendix)
7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. ☐ Computer Readable Copy
 - b. ☒ Paper Copy (identical to computer copy)
 - c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement (when there is an assignee) ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☒ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. ☐ *Small Entity Statement(s) (PTO/SB/09-12) ☐ Statement filed in prior application, Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☒ Other: 37 C.F.R. § 1.136(a)(3) Authorization
- ☒ Other: Request to Open New Disk File

*NOTE FOR ITEMS 1 & 14 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28)

17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

☐ Continuation ☒ Divisional ☐ Continuation-in-Part (CIP) of prior application No: 09/064,057

Prior application information: Examiner Whisenant, E. Group/Art Unit: 1634

18. CORRESPONDENCE ADDRESS

☐ Customer Number
or Bar Code Label

(Insert Customer No. or Attach bar code label here)

or ☒ Correspondence
address below

NAME	STERNE, KESSLER, GOLDSTEIN & FOX P L L C				
	Attorneys at Law				
ADDRESS	Suite 600, 1100 New York Avenue, NW				
CITY	Washington	STATE	DC	ZIP CODE	20005-3934
COUNTRY	USA	TELEPHONE	(202) 371-2600	FAX	(202) 371-2540

NAME (Print/Type)	Brian J. Del Buono	Registration No. (Attorney/Agent)	42,473
SIGNATURE		Date	February 5, 1999

Burden Hour Statement this form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

GERARD *et al.*

Appl. No.: (To be assigned; Divisional of
U.S. Appl. No. 09/064,057)

Filed: (Herewith)

For: **Compositions and Methods for
Reverse Transcription of Nucleic
Acid Molecules**

Art Unit: (To be assigned)

Examiner: (To be assigned)

Atty. Docket: 0942.4330003/RWE/BJD

Preliminary Amendment

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

In advance of prosecution in the above identified matter, please amend the
application as follows:

In the Specification:

Please amend the specification as follows:

At page 1, line 4, after "present application" please insert -- is a divisional of U.S.
Application No. 09/064,057, filed April 22, 1998, which --.

In the Claims:

Please cancel claims 10-21, 26-40, 46-80, and 89-115, without prejudice to or
disclaimer of the subject matter contained therein.

Please add the following new claim:

-- 116. An ASLV reverse transcriptase, wherein said ASLV reverse transcriptase is reduced or substantially reduced in RNase H activity. --

Please amend the remaining claims as follows:

41. (Once amended) An ASLV reverse transcriptase produced [according to the method of claim 26.] by a method comprising

(a) obtaining a host cell comprising one or more nucleic acid sequences encoding one or more subunits of ASLV reverse transcriptase; and

(b) culturing said host cell under conditions sufficient to produce said ASLV reverse transcriptase subunits.

86. (Once amended) The kit of claim 85, wherein said ASLV RT [is] comprises an α and a β subunit.

Remarks

No new matter has been added by these amendments. Applicants have amended the specification solely to update the priority information, and have amended the claims to cancel claims restricted in the parent application into groups other than Group I, to

introduce the limitations of canceled original claim 26 (from which claim 41 originally depended) into claim 41, and to correct an obvious typographical error in claim 86.

Support for the amendment to claim 41 may therefore be found in claim 26 as originally filed, and support for newly added claim 116 may be found throughout the specification as originally filed, particularly at page 27, lines 7-13. Hence, the foregoing amendments do not add new matter, and their entry is respectfully requested.

Upon entry of the foregoing amendments, claims 1-9, 22-25, 41-45, 81-88 and 116 are pending, with claims 1, 22, 41, 81, 85, and 116 being the independent claims.

It is respectfully believed that this application is now in condition for examination. Early notice to this effect is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Brian J. Del Buono
Agent for Applicants
Registration No. 42,473

Date: February 5, 1999

1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005
(202) 371-2600

P:\USERS\BRIAND\0942\4330003\020599.amd

Compositions and Methods for Reverse Transcription of Nucleic Acid Molecules

CROSS REFERENCE TO RELATED APPLICATIONS

5 The present application claims the benefit of U.S. Provisional Application
Nos. 60/044,589, filed April 22, 1997, and 60/049,874, filed June 17, 1997, the
disclosures of which are entirely incorporated by reference herein.

FIELD OF THE INVENTION

10 The present invention is in the fields of molecular and cellular biology.
The invention is generally related to reverse transcriptase enzymes and methods
for the reverse transcription of nucleic acid molecules, especially messenger RNA
molecules. Specifically, the invention relates to compositions comprising mixtures
of reverse transcriptase enzymes, and to methods of producing, amplifying or
sequencing nucleic acid molecules (particularly cDNA molecules) using these
reverse transcriptase enzymes or compositions. The invention also relates to
15 nucleic acid molecules produced by these methods and to the use of such nucleic
acid molecules to produce desired polypeptides. The invention also concerns kits
comprising such compositions.

BACKGROUND OF THE INVENTION

cDNA and cDNA Libraries

20 In examining the structure and physiology of an organism, tissue or cell,
it is often desirable to determine its genetic content. The genetic framework of an
organism is encoded in the double-stranded sequence of nucleotide bases in the
deoxyribonucleic acid (DNA) which is contained in the somatic and germ cells of
the organism. The genetic content of a particular segment of DNA, or gene, is
25 only manifested upon production of the protein which the gene encodes. In order

to produce a protein, a complementary copy of one strand of the DNA double helix (the "coding" strand) is produced by polymerase enzymes, resulting in a specific sequence of ribonucleic acid (RNA). This particular type of RNA, since it contains the genetic message from the DNA for production of a protein, is called messenger RNA (mRNA).

Within a given cell, tissue or organism, there exist myriad mRNA species, each encoding a separate and specific protein. This fact provides a powerful tool to investigators interested in studying genetic expression in a tissue or cell -- mRNA molecules may be isolated and further manipulated by various molecular biological techniques, thereby allowing the elucidation of the full functional genetic content of a cell, tissue or organism.

One common approach to the study of gene expression is the production of complementary DNA (cDNA) clones. In this technique, the mRNA molecules from an organism are isolated from an extract of the cells or tissues of the organism. This isolation often employs solid chromatography matrices, such as cellulose or agarose, to which oligomers of thymidine (T) have been complexed. Since the 3' termini on most eukaryotic mRNA molecules contain a string of adenosine (A) bases, and since A binds to T, the mRNA molecules can be rapidly purified from other molecules and substances in the tissue or cell extract. From these purified mRNA molecules, cDNA copies may be made using the enzyme reverse transcriptase (RT), which results in the production of single-stranded cDNA molecules. The single-stranded cDNAs may then be converted into a complete double-stranded DNA copy (*i.e.*, a double-stranded cDNA) of the original mRNA (and thus of the original double-stranded DNA sequence, encoding this mRNA, contained in the genome of the organism) by the action of a DNA polymerase. The protein-specific double-stranded cDNAs can then be inserted into a plasmid or viral vector, which is then introduced into a host bacterial, yeast, animal or plant cell. The host cells are then grown in culture

media, resulting in a population of host cells containing (or in many cases, expressing) the gene of interest.

This entire process, from isolation of mRNA to insertion of the cDNA into a plasmid or vector to growth of host cell populations containing the isolated gene, is termed "cDNA cloning." If cDNAs are prepared from a number of different mRNAs, the resulting set of cDNAs is called a "cDNA library," an appropriate term since the set of cDNAs represents a "population" of genes comprising the functional genetic information present in the source cell, tissue or organism. Genotypic analysis of these cDNA libraries can yield much information on the structure and function of the organisms from which they were derived.

Retroviral Reverse Transcriptase Enzymes

Three prototypical forms of retroviral RT have been studied thoroughly. Moloney Murine Leukemia Virus (M-MLV) RT contains a single subunit of 78 kDa with RNA-dependent DNA polymerase and RNase H activity. This enzyme has been cloned and expressed in a fully active form in *E.coli* (reviewed in Prasad, V.R., *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, p.135 (1993)). Human Immunodeficiency Virus (HIV) RT is a heterodimer of p66 and p51 subunits in which the smaller subunit is derived from the larger by proteolytic cleavage. The p66 subunit has both a RNA-dependent DNA polymerase and an RNase H domain, while the p51 subunit has only a DNA polymerase domain. Active HIV p66/p51 RT has been cloned and expressed successfully in a number of expression hosts, including *E.coli* (reviewed in Le Grice, S.F.J., *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory press, p. 163 (1993)). Within the HIV p66/p51 heterodimer, the 51-kD subunit is catalytically inactive, and the 66-kD subunit has both DNA polymerase and RNase H activity (Le Grice, S.F.J., *et al.*, *EMBO Journal* 10:3905 (1991); Hostomsky, Z., *et al.*, *J. Virol.* 66:3179 (1992)). Avian Sarcoma-Leukosis Virus (ASLV) RT, which includes but is not limited to

Rous Sarcoma Virus (RSV) RT, Avian Myeloblastosis Virus (AMV) RT, Avian Erythroblastosis Virus (AEV) Helper Virus MCAV RT, Avian Myelocytomatosis Virus MC29 Helper Virus MCAV RT, Avian Reticuloendotheliosis Virus (REV-T) Helper Virus REV-A RT, Avian Sarcoma Virus UR2 Helper Virus UR2AV RT, Avian Sarcoma Virus Y73 Helper Virus YAV RT, Rous Associated Virus (RAV) RT, and Myeloblastosis Associated Virus (MAV) RT, is also a heterodimer of two subunits, α (approximately 62 kDa) and β (approximately 94 kDa), in which α is derived from β by proteolytic cleavage (reviewed in Prasad, V.R., *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1993), p. 135). ASLV RT can exist in two additional catalytically active structural forms, $\beta\beta$ and $\alpha\beta$ (Hizi, A. and Joklik, W.K., *J. Biol. Chem.* 252: 2281 (1977)). Sedimentation analysis suggests $\alpha\beta$ and $\beta\beta$ are dimers and that the α form exists in an equilibrium between monomeric and dimeric forms (Grandgenett, D.P., *et al.*, *Proc. Nat. Acad. Sci. USA* 70: 230 (1973); Hizi, A. and Joklik, W.K., *J. Biol. Chem.* 252: 2281 (1977); and Soltis, D.A. and Skalka, A.M., *Proc. Nat. Acad. Sci. USA* 85: 3372 (1988)). The ASLV $\alpha\beta$ and $\beta\beta$ RTs are the only known examples of retroviral RT that include three different activities in the same protein complex: DNA polymerase, RNase H, and DNA endonuclease (integrase) activities (reviewed in Skalka, A.M., *Reverse Transcriptase*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1993), p. 193). The α form lacks the integrase domain and activity.

Various forms of the individual subunits of ASLV RT have been cloned and expressed. These include a 98-kDa precursor polypeptide that is normally processed proteolytically to β and a 4-kDa polypeptide removed from the β carboxy end (Alexander, F., *et al.*, *J. Virol.* 61: 534 (1987) and Anderson, D. *et al.*, *Focus* 17:53 (1995)), and the mature β subunit (Weis, J.H. and Salstrom, J.S., U.S. Patent No. 4, 663, 290 (1987); and Soltis, D.A. and Skalka, A.M., *Proc. Nat. Acad. Sci. USA* 85:3372 (1988)). Heterodimeric RSV $\alpha\beta$ RT has also been purified from *E. coli* cells expressing a cloned RSV β gene (Chernov, A.P., *et al.*,

Biomed. Sci. 2:49 (1991)). However, there have been no reports heretofore of the simultaneous expression of cloned ASLV RT α and β genes resulting in the formation of heterodimeric $\alpha\beta$ RT.

Reverse Transcription Efficiency

As noted above, the conversion of mRNA into cDNA by RT-mediated reverse transcription is an essential step in the study of proteins expressed from cloned genes. However, the use of unmodified RT to catalyze reverse transcription is inefficient for at least two reasons. First, RT sometimes destroys an RNA template before reverse transcription is initiated, primarily due to the activity of intrinsic RNase H activity present in RT. Second, RT often fails to complete reverse transcription after the process has been initiated (Berger, S.L., *et al.*, *Biochemistry* 22:2365-2372 (1983); Krug, M.S., and Berger, S.L., *Meth. Enzymol.* 152:316 (1987)). Removal of the RNase H activity of RT can eliminate the first problem and improve the efficiency of reverse transcription (Gerard, G.F., *et al.*, *FOCUS* 11(4):60 (1989); Gerard, G.F., *et al.*, *FOCUS* 14(3):91 (1992)). However RTs, including those forms lacking RNase H activity ("RNase H" forms), still tend to terminate DNA synthesis prematurely at certain secondary structural (Gerard, G.F., *et al.*, *FOCUS* 11(4):60 (1989); Myers, T.W., and Gelfand, D.H., *Biochemistry* 30:7661 (1991)) and sequence (Messer, L.I., *et al.*, *Virol.* 146:146 (1985)); Abbotts, J., *et al.*, *J. Biol. Chem.* 268:10312-10323 (1993)) barriers in nucleic acid templates.

Even in the most efficient reverse transcription systems available today, which use RNase H⁻ M-MLV RT, yields of total cDNA product generally do not exceed 50% of input mRNA and the fraction of the product that is full-length does not exceed 50%. The secondary structural and sequence barriers in the mRNA template, which as described above can give rise to these limitations, occur frequently at homopolymer stretches (Messer, L.I., *et al.*, *Virol.* 146:146 (1985); Huber, H.E., *et al.*, *J. Biol. Chem.* 264:4669-4678 (1989); Myers, T.W., and

Gelfand, D.H., *Biochemistry* 30:7661 (1991)), are more often sequence rather than secondary structural barriers (Abbotts, J., *et al.*, *J. Biol. Chem.* 268:10312-10323 (1993)), and are often distinct for different RTs (Abbotts, J., *et al.*, *J. Biol. Chem.* 268:10312-10323 (1993)). If these barriers could be overcome, yield of total and full-length cDNA product in reverse transcription reactions could be increased.

SUMMARY OF THE INVENTION

The present invention provides reverse transcriptase enzymes, compositions comprising such enzymes and methods useful in overcoming the above-described cDNA length limitations. In general, the invention provides compositions for use in reverse transcription of a nucleic acid molecule comprising two or more different polypeptides having reverse transcriptase activity. Such compositions may further comprise one or more nucleotides, a suitable buffer, and/or one or more DNA polymerases. The compositions of the invention may also comprise one or more oligonucleotide primers. Each reverse transcriptase used in the compositions of the invention may have a different transcription pause site on a given mRNA molecule. The reverse transcriptases in these compositions preferably are reduced or substantially reduced in RNase H activity, and most preferably are enzymes selected from the group consisting of Moloney Murine Leukemia Virus (M-MLV) H⁻ reverse transcriptase, Rous Sarcoma Virus (RSV) H⁻ reverse transcriptase, Avian Myeloblastosis Virus (AMV) H⁻ reverse transcriptase, Rous Associated Virus (RAV) H⁻ reverse transcriptase, Myeloblastosis Associated Virus (MAV) H⁻ reverse transcriptase and Human Immunodeficiency Virus (HIV) H⁻ reverse transcriptase or other ASLV H⁻ reverse transcriptases. In preferred compositions, the reverse transcriptases are present at working concentrations.

5 The invention is also directed to methods for making one or more nucleic acid molecules, comprising mixing one or more nucleic acid templates (preferably one or more RNA templates and most preferably one or more messenger RNA templates) with two or more polypeptides having reverse transcriptase activity and incubating the mixture under conditions sufficient to make a first nucleic acid molecule or molecules complementary to all or a portion of the one or more nucleic acid templates. In a preferred embodiment, the first nucleic acid molecule is a single-stranded cDNA. Nucleic acid templates suitable for reverse transcription according to this aspect of the invention include any nucleic acid molecule or population of nucleic acid molecules (preferably RNA and most preferably mRNA), particularly those derived from a cell or tissue. In a preferred aspect, a population of mRNA molecules (a number of different mRNA molecules, typically obtained from cells or tissue) are used to make a cDNA library, in accordance with the invention. Preferred cellular sources of nucleic acid templates include bacterial cells, fungal cells, plant cells and animal cells.

10 The invention also concerns methods for making one or more double-stranded nucleic acid molecules. Such methods comprise (a) mixing one or more nucleic acid templates (preferably RNA or mRNA, and more preferably a population of mRNA templates) with two or more polypeptides having reverse transcriptase activity; (b) incubating the mixture under conditions sufficient to make a first nucleic acid molecule or molecules complementary to all or a portion of the one or more templates; and (c) incubating the first nucleic acid molecule under conditions sufficient to make a second nucleic acid molecule or molecules complementary to all or a portion of the first nucleic acid molecule or molecules, thereby forming one or more double-stranded nucleic acid molecules comprising the first and second nucleic acid molecules. Such methods may include the use of one or more DNA polymerases as part of the process of making the one or more double-stranded nucleic acid molecules. The invention also concerns compositions useful for making such double-stranded nucleic acid molecules.

Such compositions comprise two or more reverse transcriptases and optionally one or more DNA polymerases, a suitable buffer and one or more nucleotides.

5 The invention also relates to methods for amplifying a nucleic acid molecule. Such amplification methods comprise mixing the double-stranded nucleic acid molecule or molecules produced as described above with one or more DNA polymerases and incubating the mixture under conditions sufficient to amplify the double-stranded nucleic acid molecule. In a first preferred embodiment, the invention concerns a method for amplifying a nucleic acid molecule, the method comprising (a) mixing one or more nucleic acid templates (preferably one or more RNA or mRNA templates and more preferably a population of mRNA templates) with two or more different polypeptides having reverse transcriptase activity and with one or more DNA polymerases and (b) incubating the mixture under conditions sufficient to amplify nucleic acid molecules complementary to all or a portion of the one or more templates. Preferably, the reverse transcriptases are reduced or substantially reduced in RNase H activity and the DNA polymerases comprise a first DNA polymerase having 3' exonuclease activity and a second DNA polymerase having substantially reduced 3' exonuclease activity. The invention also concerns compositions comprising two or more reverse transcriptases and one or more DNA polymerases for use in amplification reactions. Such compositions may further comprise one or more nucleotides and a buffer suitable for amplification. The compositions of the invention may also comprise one or more oligonucleotide primers.

20 In accordance with the invention, at least two, at least three, at least four, at least five, at least six, or more, reverse transcriptases may be used. Preferably, 25 two to six, two to five, two to four, two to three, and most preferably two, reverse transcriptases are used in the compositions and methods of the invention. Such multiple reverse transcriptases may be added simultaneously or sequentially in any order to the compositions or in the methods of the invention. Alternatively, multiple different reactions with different enzymes may be performed separately

and the reaction products may be mixed. Thus, the invention relates to the synthesis of the nucleic acid molecules by the methods of the invention in which multiple reverse transcriptases are used simultaneously or sequentially or separately. In particular, the invention relates to a method of making one or more nucleic acid molecules comprising incubating one or more nucleic acid templates (preferably one or more RNA templates or mRNA templates, and more preferably a population of mRNA templates) with a first reverse transcriptase under conditions sufficient to make one or more nucleic acid molecules complementary to all or a portion of the one or more templates. In accordance with the invention, the one or more nucleic acid molecules (including mRNA templates and/or synthesized nucleic acid molecules) may be incubated with a second reverse transcriptase under conditions sufficient to make additional nucleic acid molecules complementary to all or a portion of the templates or to increase the length of the previously made nucleic acid molecules. In accordance with the invention, this procedure may be repeated any number of times with the same or different reverse transcriptases of the invention. For example, the first and second reverse transcriptases may be the same or different. Furthermore, the first and third reverse transcriptases (in aspects of the invention where the procedure is repeated three times using a first, second, and third reverse transcriptase) may be the same while the second reverse transcriptase may be different from the first and the third reverse transcriptase. Thus, any combination of the same and/or different reverse transcriptases may be used in this aspect of the invention. Preferably, when multiple reverse transcriptases are used, at least two reverse transcriptases are different.

In a related aspect of the invention, the reverse transcriptase used in the reaction may retain all or a portion of its activity during subsequent reaction steps. Alternatively, the reverse transcriptase used in the reaction may be inactivated by any method prior to incubation with additional reverse transcriptases. Such an

inactivation may include but is not limited to heat inactivation, organic extraction (*e.g.*, with phenol and/or chloroform), ethanol precipitation and the like.

The synthesized nucleic acid molecules made by simultaneous or sequential or separate addition of reverse transcriptases may then be used to make double stranded nucleic acid molecules. Such synthesized nucleic acid molecules serve as a template which when incubated under appropriate conditions (*e.g.*, preferably in the presence of one or more DNA polymerases) make nucleic acid molecules complementary to all or a portion of the synthesized nucleic acid molecules, thereby forming a number of double stranded nucleic acid molecules. The double stranded molecules may then be amplified in accordance with the invention.

The invention is also directed to nucleic acid molecules (particularly single- or double-stranded cDNA molecules) or amplified nucleic acid molecules produced according to the above-described methods and to vectors (particularly expression vectors) comprising these nucleic acid molecules or amplified nucleic acid molecules.

The invention is also directed to recombinant host cells comprising the above-described nucleic acid molecules, amplified nucleic acid molecules or vectors. Preferred such host cells include bacterial cells, yeast cells, plant cells and animal cells (including insect cells and mammalian cells).

The invention is further directed to methods of producing a polypeptide comprising culturing the above-described recombinant host cells and isolating the polypeptide, and to a polypeptide produced by such methods.

The invention also concerns methods for sequencing one or more nucleic acid molecules using the compositions or enzymes of the invention. Such methods comprise (a) mixing one or more nucleic acid molecules (*e.g.*, one or more RNA or DNA molecules) to be sequenced with one or more primers, one or more polypeptides having reverse transcriptase activity, one or more nucleotides and one or more terminating agents, such as one or more dideoxynucleoside triphosphates; (b) incubating the mixture under conditions sufficient to synthesize

a population of nucleic acid molecules complementary to all or a portion of the one or more nucleic acid molecules to be sequenced; and (c) separating the population of nucleic acid molecules to determine the nucleotide sequence of all or a portion of the one or more nucleic acid molecules to be sequenced. In these sequencing methods of the invention, the one or more polypeptides having reverse transcriptase activity may be added simultaneously, sequentially, or separately to the reaction mixtures as described above.

The invention is also directed to kits for use in the methods of the invention. Such kits can be used for making, sequencing or amplifying nucleic acid molecules (single- or double-stranded). The kits of the invention comprise a carrier, such as a box or carton, having in close confinement therein one or more containers, such as vials, tubes, bottles and the like. In the kits of the invention, a first container contains one or more of the reverse transcriptase enzymes (preferably one or more such enzymes that are reduced or substantially reduced in RNase H activity) or one or more of the compositions of the invention. In another aspect, the kit may contain one or more containers comprising two or more, three or more, four or more, five or more, six or more, and the like, reverse transcriptases, preferably one or more containers comprising two to six, two to five, two to four, two to three, or more preferably two, reverse transcriptases. The kits of the invention may also comprise, in the same or different containers, one or more DNA polymerase (preferably thermostable DNA polymerases), a suitable buffer for nucleic acid synthesis and one or more nucleotides. Alternatively, the components of the composition may be divided into separate containers (*e.g.*, one container for each enzyme). In preferred kits of the invention, the reverse transcriptases are reduced or substantially reduced in RNase H activity, and are most preferably selected from the group consisting of M-MLV H⁻ reverse transcriptase, RSV H⁻ reverse transcriptase, AMV H⁻ reverse transcriptase, RAV H⁻ reverse transcriptase, MAV H⁻ reverse transcriptase and HIV H⁻ reverse transcriptase. In additional preferred kits of the invention, the

enzymes (reverse transcriptases and/or DNA polymerases) in the containers are present at working concentrations.

The invention also relates to methods of producing RSV reverse transcriptase (and/or subunits thereof). In particular, the invention relates to methods for producing RSV reverse transcriptase (and/or subunits thereof) containing RNase H activity, to methods for producing RSV reverse transcriptase (and/or subunits thereof) that is reduced or substantially reduced in RNase H activity, and to RSV reverse transcriptases (and/or subunits thereof) produced by such methods.

The invention further relates to methods for using such reverse transcriptases and to kits comprising such reverse transcriptases. In particular, the RSV reverse transcriptases (and/or subunits thereof) of the invention may be used in methods of sequencing, amplification and production (via, *e.g.*, reverse transcription) of nucleic acid molecules.

The invention also relates to methods of producing AMV reverse transcriptase (and/or subunits thereof). In particular, the invention relates to methods for producing AMV reverse transcriptase (and/or subunits thereof) containing RNase H activity, to methods for producing AMV reverse transcriptase (and/or subunits thereof) that is reduced or substantially reduced in RNase H activity, and to AMV reverse transcriptases (and/or subunits thereof) produced by such methods.

The invention further relates to methods for using such reverse transcriptases and to kits comprising such reverse transcriptases. In particular, the AMV reverse transcriptases (and/or subunits thereof) of the invention may be used in methods of sequencing, amplification and production (via, *e.g.*, reverse transcription) of nucleic acid molecules.

The invention also generally relates to methods of producing ASLV reverse transcriptases (and/or subunits thereof). In particular, the invention relates to methods for producing ASLV reverse transcriptases (and/or subunits thereof)

containing RNase H activity, to methods for producing such ASLV reverse transcriptases that are reduced or substantially reduced in RNase H activity, and to ASLV reverse transcriptases produced by such methods.

5 The invention further relates to methods for using such reverse transcriptases and to kits comprising such reverse transcriptases. In particular, the ASLV reverse transcriptases (and/or subunits thereof) of the invention may be used in methods of sequencing, amplification and production (*e.g.*, via reverse transcription) of nucleic acid molecules.

10 The invention further relates to methods for elevated- or high-temperature reverse transcription of a nucleic acid molecule comprising (a) mixing one or more nucleic acid templates (preferably one or more RNA molecules (*e.g.*, one or more mRNA molecules or polyA⁺ RNA molecules, and more preferably a population of mRNA molecules) or one or more DNA molecules) with one or more polypeptides having reverse transcriptase activity; and (b) incubating the mixture at a temperature of 50°C or greater and under conditions sufficient to make a first
15 nucleic acid molecule or molecules (such as a full length cDNA molecule) complementary to all or a portion of the one or more nucleic acid templates. In a preferred aspect, a population of mRNA molecules is used to make a cDNA library at elevated or high temperatures. In another aspect, elevated- or high-temperature nucleic acid synthesis is conducted with multiple reverse
20 transcriptases (*i.e.*, two or more, three or more, four or more, five or more, six or more, and the like, more preferably two to six, two to five, two to four, two to three, and still more preferably two, reverse transcriptases), which may be added to the reaction mixture simultaneously or sequentially or separately as described above. In preferred such methods, the mixture is incubated at a temperature of
25 about 51°C or greater, about 52°C or greater, about 53°C or greater, about 54°C or greater, about 55°C or greater, about 56°C or greater, about 57°C or greater, about 58°C or greater, about 59°C or greater, about 60°C or greater, about 61°C or greater, about 62°C or greater, about 63°C or greater, about 64°C or greater,

about 65°C or greater, about 66°C or greater, about 67°C or greater, about 68°C or greater, about 69°C or greater, about 70°C or greater, about 71°C or greater, about 72°C or greater, about 73°C or greater, about 74°C or greater, about 75°C or greater, about 76°C or greater, about 77°C or greater, or about 78°C or greater; or at a temperature range of from about 50°C to about 75°C, about 51°C to about 75°C, about 52°C to about 75°C, about 53°C to about 75°C, about 54°C to about 75°C, about 55°C to about 75°C, about 50°C to about 70°C, about 51°C to about 70°C, about 52°C to about 70°C, about 53°C to about 70°C, about 54°C to about 70°C, about 55°C to about 70°C, about 55°C to about 65°C, about 56°C to about 65°C, about 56°C to about 64°C or about 56°C to about 62°C. The invention is also directed to such methods which further comprise incubating the first nucleic acid molecule or molecules under conditions sufficient to make a second nucleic acid molecule or molecules complementary to all or portion of the first nucleic acid molecule or molecules. According to the invention, the first and second nucleic acid molecules produced by these methods may be DNA molecules, and may form a double stranded DNA molecule or molecules which may be a full length cDNA molecule or molecules, such as a cDNA library. The one or more polypeptides having reverse transcriptase activity that are used in these methods preferably are reduced or substantially reduced in RNase H activity, and are preferably selected from ASLV reverse transcriptases (and/or subunits thereof) such as one or more subunits of AMV reverse transcriptase and/or one or more subunits of RSV reverse transcriptase and/or one or more subunits of MAV reverse transcriptase, and/or one or more subunits of RAV reverse transcriptase, particularly wherein the subunits are reduced or substantially reduced in RNase H activity.

The invention also relates to kits for elevated- or high-temperature nucleic acid synthesis, which may comprise one or more components selected from the group consisting of one or more reverse transcriptases (preferably one or more ASLV reverse transcriptases such as AMV or RSV reverse transcriptases (or one

or more subunits thereof), and more preferably one or more AMV or RSV reverse transcriptases (or one or more subunits thereof) which are reduced or substantially reduced in RNase H activity), one or more nucleotides, one or more primers and one or more suitable buffers.

5 The invention is also directed to nucleic acid molecules produced by the above-described methods which may be full-length cDNA molecules, to vectors (particularly expression vectors) comprising these nucleic acid molecules and to host cells comprising these vectors and nucleic acid molecules.

10 Other preferred embodiments of the present invention will be apparent to one of ordinary skill in light of the following drawings and description of the invention, and of the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

15 **Figures 1-6** describe, in schematic form (with details omitted for clarity), the construction of the expression vector (pDABH^{His}) which places the RSV RT α and β genes under control of insect viral promoters:

Figure 1: pJD100, pAMP18, pAMP18N, pAMP1, pAMP1C, pAMP18NM and pAMP18B.

Figure 2: M13, M13RT, pAMP18BH- and M13RTH-.

Figure 3: pAMP1A.

20 **Figure 4:** pDBH-Kpn, pDABH-, pDBH-KpnHis.

Figure 5: pFastBac DUAL, pFastBac DUAL Nde, pDBH- and pDA.

Figure 6: pDABH-His.

Figures 7-21 are more detailed maps of the plasmids described in Figures 1-6:

25 **Figure 7:** pJD100.

Figure 8: pAMP18N.

Figure 9: pAMP1C.

Figure 10: pAMP18NM.

Figure 11: pAMP18B.

Figure 12: M13RT.

Figure 13: M13RTH-.

Figure 14: pAMP18BH-.

Figure 15: pDBH-.

Figure 16: pAMP1A.

Figure 17: pDBH-Kpn.

Figure 18: pDBH-KpnHis.

Figure 19: pDA.

Figure 20: pDABH-.

Figure 21: pDABH-His.

Figures 22-25 describe, in schematic form (with details omitted for clarity), the construction of the expression vector (pDAMVAH-BH-) which places the AMV RT α and β genes under control of insect viral promoters:

Figure 22: Cloning of AMV RT gene from RNA; pSPORT8.

Figure 23: Construction of a His-tagged AMV RT β gene; pAMVN, pAMVNM, pAMVNMH-, pAMVC and pAMVBH-.

Figure 24: Construction of clones for the AMV RT α subunit by PCR; pAMVA and pAMVAH-.

Figure 25: Construction of vectors comprising the AMV RT α and β genes; pD, pDAMVAH-, pDAMVA, pAMVH-BH-, pDAMVABH- and pJAMVBH-.

Figures 26-38 are more detailed maps of the plasmids described in Figures 22-25:

Figure 26: pAMVN.

Figure 27: pAMVC.

Figure 28: pAMVNM.

Figure 29: pAMVNMH-.

Figure 30: pAMVBH-.

Figure 31: pAMVA.

Figure 32: pAMVAH-.

Figure 33: pFastBacDual (pD).

Figure 34: pDAMVA.

Figure 35: pDAMVAH-.

Figure 36: pDAMVABH-.

Figure 37: pJAMVBH-.

Figure 38: pDAMVAH-BH-.

Figure 39 is a semi-logarithmic graph demonstrating RT activities of various RTs incubated for the times and at the temperatures indicated.

Figure 40 is an autoradiograph of cDNA products synthesized by various RTs at the temperatures (°C) indicated from 1.4-, 2.4-, 4.4- and 7.5-Kb mRNAs over 50-minute reactions. M: ³²P-labeled 1Kb DNA ladder.

Figure 41 is an autoradiograph of cDNA products synthesized by RSV H⁻ RT and SS II RT at the temperatures (°C) indicated from 1.4-, 2.4-, 4.4-, and 7.5-Kb mRNAs over 30-minute reactions. M: ³²P-labeled 1 Kb DNA ladder.

Figure 42 is a graph of the amounts of full length cDNA synthesized by SSII and RSV H⁻ RT in Figure 41 as a function of incubation temperature.

Figure 43 is a restriction map of plasmid pBP-RT(PCR).

Figure 44 is a restriction map of plasmid pBP-RT(ATG).

Figure 45 is a restriction map of plasmid pBK-RT15(ATG).

Figure 46 is a restriction map of plasmid pFBBH-His.

Figure 47 is a restriction map of plasmid pJB-His.

Figure 48 is a restriction map of plasmid pJBD110E-His.

Figure 49 is a restriction map of plasmid pDAD110E.

Figure 50 is a restriction map of plasmid pFBBD110E-His.

Figure 51 is a restriction map of plasmid pDABHis.

Figure 52 is a restriction map of plasmid pDABD110EHis.

Figure 53 is a restriction map of plasmid pDAD110EBHis.

Figure 54 is a restriction map of plasmid pDAD110EB110E.

DETAILED DESCRIPTION OF THE INVENTION

Overview

5 The present invention provides compositions and methods useful in overcoming the length limitations often observed during reverse transcription of nucleic acid molecules. Thus, the invention facilitates the production of full-length cDNA molecules not heretofore possible.

10 In general, the invention provides compositions for use in reverse transcription of a nucleic acid molecule comprising two or more, three or more, four or more, five or more, six or more, and the like, different polypeptides having reverse transcriptase activity. The compositions of the invention preferably comprise two to six, two to five, two to four, two to three, and more preferably
15 comprise two, polypeptides having reverse transcriptase activity. The enzymes in these compositions are preferably present in working concentrations and are reduced or substantially reduced in RNase H activity, although mixtures of enzymes, some having RNase H activity and some reduced or substantially reduced in RNase H activity, may be used in the compositions of the invention. Alternatively, the reverse transcriptases used in the compositions of the invention
20 may have RNase H activity. Preferred reverse transcriptases include M-MLV H⁻ reverse transcriptase, RSV H⁻ reverse transcriptase, AMV H⁻ reverse transcriptase, RAV H⁻ reverse transcriptase, MAV H⁻ reverse transcriptase and HIV H⁻ reverse transcriptase or other ASLV H⁻ reverse transcriptases.

25 The invention is also directed to methods for reverse transcription of one or more nucleic acid molecules comprising mixing one or more nucleic acid templates, which is preferably RNA or messenger RNA (mRNA) and more

preferably a population of mRNA molecules, with two or more polypeptides having reverse transcriptase activity (or with the compositions of the invention) and incubating the mixture under conditions sufficient to make a nucleic acid molecule or molecules complementary to all or a portion of the one or more templates. Such nucleic acid synthesis may be accomplished by sequential or simultaneous or separate addition of multiple reverse transcriptases. Preferably, two or more, three or more, four or more, five or more, six or more, and the like, reverse transcriptases are used, or a range of two to six, two to five, two to four, two to three and more preferably two, reverse transcriptases are used. To make the nucleic acid molecule or molecules complementary to the one or more templates, a primer (*e.g.*, an oligo(dT) primer) and one or more nucleotides are used for nucleic acid synthesis in the 3' to 5' direction. Nucleic acid molecules suitable for reverse transcription according to this aspect of the invention include any nucleic acid molecule, particularly those derived from a prokaryotic or eukaryotic cell. Such cells may include normal cells, diseased cells, transformed cells, established cells, progenitor cells, precursor cells, fetal cells, embryonic cells, bacterial cells, yeast cells, animal cells (including human cells), avian cells, plant cells and the like, or tissue isolated from a plant or an animal (*e.g.*, human, cow, pig, mouse, sheep, horse, monkey, canine, feline, rat, rabbit, bird, fish, insect, etc.). Such nucleic acid molecules may also be isolated from viruses.

The invention further provides methods for amplifying or sequencing a nucleic acid molecule comprising contacting the nucleic acid molecule with two or more polypeptides having reverse transcriptase activity (or with the compositions of the invention). Such reactions may be accomplished by sequential or simultaneous or separate addition of the two or more polypeptides having reverse transcriptase activity to the reaction mixtures. Preferred such methods comprise one or more polymerase chain reactions (PCRs).

The invention also provides cDNA molecules or amplified nucleic acid molecules produced according to the above-described methods, vectors

(particularly expression vectors) comprising these cDNA molecules or amplified nucleic acid molecules, and recombinant host cells comprising such cDNA molecules, amplified nucleic acid molecules or vectors. The invention also provides methods of producing a polypeptide comprising culturing these recombinant host cells and isolating the polypeptide, and provides a polypeptide produced by such methods.

The invention also provides kits for use in accordance with the invention. Such kits comprise a carrier means, such as a box or carton, having in close confinement therein one or more container means, such as vials, tubes, bottles and the like, wherein the kit comprises, in the same or different containers, two or more polypeptides having reverse transcriptase activity. The kits of the invention may also comprise, in the same or different containers, one or more DNA polymerases, a suitable buffer and/or one or more nucleotides (such as deoxynucleoside triphosphates (dNTPs)).

The invention also concerns a substantially pure RSV reverse transcriptase (RSV RT), which may or may not be reduced or substantially reduced in RNase H activity. Such RSV RTs may comprise one or more subunits (or derivatives, variants, fragments or mutants thereof) selected from one or more α subunits, one or more β subunits, and one or more $\beta p4$ subunits, any or all of which may or may not be reduced or substantially reduced in RNase H activity. In one preferred aspect of the invention, the RSV RT may comprise an α subunit reduced or substantially reduced in RNase H activity and a β subunit having RNase H activity (*i.e.*, RSV $\alpha H^- \beta H^+$ RT). In a preferred aspect of this embodiment, the gene encoding the α subunit has been modified or mutated to reduce RNase H activity while the gene encoding the β subunit has not been mutated or modified in this manner. Such mutations or modifications are preferably made within the RNase H domain of the α subunit. Unexpectedly, the phenotype of this construct showed substantially reduced (*i.e.*, approximately 5% of wildtype) RNase H activity. In another preferred aspect, the RSV RT may comprise an α subunit reduced or

substantially reduced in RNase H activity and a β subunit also reduced or substantially reduced in RNase H activity (*i.e.*, RSV $\alpha H^-/\beta H^-$ RT). In another preferred aspect, the RSV RT may comprise two β subunits, either or both of which may or may not be reduced or substantially reduced in RNase H activity (*i.e.*, RSV $\beta H^-/\beta H^-$ RT; RSV $\beta H^-/\beta H^+$ RT; or RSV $\beta H^+/\beta H^+$ RT). In another preferred aspect, the RSV RT may comprise a single α subunit, which may or may not be reduced or substantially reduced in RNase H activity (*i.e.*, RSV αH^- RT or RSV αH^+ RT). In another preferred aspect, the RSV RT may comprise one or more $\beta p4$ subunits, any or all of which may or may not be reduced or substantially reduced in RNase H activity (*e.g.*, RSV $\beta p4 H^-/\beta p4 H^-$ RT; RSV $\beta p4 H^-/\beta p4 H^+$ RT; RSV $\beta p4 H^+/\beta p4 H^+$ RT; RSV $\alpha H^-/\beta p4 H^+$ RT; RSV $\alpha H^+/\beta p4 H^+$ RT; RSV $\alpha H^-/\beta p4 H^-$ RT; RSV $\alpha H^+/\beta p4 H^-$ RT; RSV $\beta H^-/\beta p4 H^+$ RT; RSV $\beta H^+/\beta p4 H^+$ RT; RSV $\beta H^-/\beta p4 H^-$ RT; RSV $\beta H^+/\beta p4 H^-$ RT; etc.). As will be recognized, derivatives, variants, fragments or mutants of any or all of the above subunits may also be used in accordance with the invention.

In a related aspect of the invention, where the RSV RTs comprise two or more subunits (or derivatives, variants, fragments or mutants thereof) and preferably comprise two subunits (*e.g.*, a dimer), at least one but preferably not all of these subunits may be modified or mutated to reduce, substantially reduce or eliminate the polymerase activity of at least one subunit (*e.g.*, pol-). In a preferred aspect, for an RSV RT which comprises an α and β subunit, the β subunit has been modified or mutated (preferably by recombinant techniques) to reduce, substantially reduce or eliminate the polymerase activity while the polymerase activity of the α subunit has not been mutated or modified in this manner. Preferably, the α subunit of such RSV RT has also been modified or mutated to reduce or substantially reduce RNase H activity while the β subunit has not been mutated or modified in this manner. Such a construct may be designated $\alpha H^-/\beta H^+$ pol-. Any number of combinations of subunits can be prepared in which mutations or modifications are made in one subunit of a two subunit enzyme, and

these constructs may be combined with other modifications or mutations, such as those which reduce or substantially reduce RNase H activity. Illustrated examples include but are not limited to RSV α H-/ β H- pol-; RSV α H+/ β H+ pol-; RSV α H- pol-/ β H+; RSV α H- pol-/ β H-; RSV α H+ pol-/ β H+; RSV β H-/ β H- pol-; RSV β H-/ β H+ pol-; RSV β H+/ β H+ pol-; RSV β p4H-/ β p4H- pol-; RSV β p4H-/ β p4H+ pol-; RSV β p4H+/ β p4H+ pol-; RSV α H-/ β p4H- pol-; RSV α H+/ β p4H+ pol-; RSV α H+/ β p4H- pol-; RSV β H-/ β p4H+ pol-; RSV β H-/ β p4H- pol-; RSV β H+/ β p4H- pol-; etc. In a preferred aspect, the polymerase domain of the subunit is modified or mutated (one or more point mutations, deletion mutations and/or insertion mutations) by recombinant techniques. In another aspect, the nucleotide binding site of the polymerase domain is modified or mutated. In a preferred aspect, one or more acidic amino acids within the nucleotide binding site are substituted with different amino acids. Particularly preferred amino acids within the nucleotide binding site for mutation or modification include, but are not limited to, Asp¹⁰⁷, Leu¹⁰⁸, Lys¹⁰⁹, and Asp¹¹⁰, or the corresponding amino acid sequence.

The invention also relates to methods of producing the RSV RTs of the invention, which methods comprise obtaining a host cell comprising a nucleic acid sequence encoding one or more α subunits (or derivatives, variants, fragments or mutants thereof) and/or a nucleic acid sequence encoding one or more β subunits (or derivatives, variants, fragments or mutants thereof) and/or a nucleic acid sequence encoding one or more β p4 subunits (or derivatives, variants, fragments or mutants thereof), and culturing the host cell under conditions sufficient to produce the RSV RTs of the invention. The nucleic acid sequences encoding such α subunit(s) and/or such β subunit(s) and/or such β p4 subunit(s) may be contained in the same vector or in different vectors. In accordance with the invention, such α subunit(s) and/or β subunit(s) and/or β p4 subunit(s) may be produced separately and mixed before or after isolation of each subunit to form the RSV RTs of the invention. Alternatively, such α and/or β subunits and/or β p4 subunits may be

expressed simultaneously (*i.e.*, co-expressed) in the same host cell, thereby producing an RSV RT comprising an α and a β subunit, an α subunit alone, a β subunit alone, a β p4 subunit alone, a β subunit and a β p4 subunit, two β subunits, or two β p4 subunits, or derivatives, variants, fragments or mutants thereof. In a preferred aspect, the α subunit (or derivatives, variants, fragments or mutants thereof) is simultaneously expressed in a host cell with the β subunit (or derivatives, variants, fragments or mutants thereof). In a related aspect of the invention, the β or β p4 subunits (or derivatives, fragments or mutants thereof) may be expressed in a host or host cell which accomplishes *in vivo* processing of some or all of such β or β p4 subunits to form the corresponding α subunit. The presence of both the β and the α subunits allows *in vivo* formation of an RSV RT which comprises an α and a β subunit. Such *in vivo* processing is preferably accomplished by expressing the β and/or β p4 subunits (or derivatives, variants, fragments or mutants thereof) in a host cell, which may be prokaryotic or eukaryotic, having appropriate processing enzymes or proteins which cleave such β or β p4 subunits to form a corresponding α subunit. Such processing enzymes or proteins may be introduced and expressed in the host system by recombinant means or may exist naturally in the host system. Preferred hosts for *in vivo* processing include eukaryotic cells or organelles such as yeast, fungi, plants, animals, insects, fish, and the like. Recombinant systems (vectors, expression vectors, promoters, etc.) which allow cloning of the β or β p4 subunits (or derivatives, fragments or mutants thereof) for *in vivo* processing are well known to one of ordinary skill in the art. As noted above, any or all of the α and/or β and/or β p4 subunits of the RSV RT (or derivatives, variants, fragments or mutants thereof) produced by these recombinant techniques may be reduced or substantially reduced in RNase H activity.

These RSV RTs or subunits thereof may then be isolated from the host cell, and may be substantially purified by any method of protein purification that will be familiar to those of ordinary skill in the art (*e.g.*, chromatography,

electrophoresis, dialysis, high-salt precipitation, or combinations thereof). The invention also relates to kits comprising one or more of the RSV RTs of the invention.

The invention also concerns a substantially pure Avian Myeloblastosis Virus reverse transcriptase (AMV RT), which may or may not be reduced or substantially reduced in RNase H activity. Such AMV RTs may comprise one or more subunits selected from one or more α subunits, one or more β subunits, and one or more β p4 subunits (or derivatives, variants, fragments or mutants thereof), any or all of which may or may not be reduced or substantially reduced in RNase H activity. In one preferred aspect of the invention, the AMV RT may comprise an α subunit reduced or substantially reduced in RNase H activity and a β subunit having RNase H activity (*i.e.*, AMV α H⁻ β H⁺ RT). In a particularly preferred aspect of this embodiment, the gene encoding the α subunit has been modified or mutated to reduce RNase activity (preferably within the RNase H domain) while the gene encoding the β subunit has not been modified or mutated to affect RNase H activity. Unexpectedly, this construct demonstrates a phenotype in which the RNase H activity of the AMV RT comprising the α subunit and the β subunit is substantially reduced in RNase H activity (*i.e.*, approximately 5% of wildtype). In another preferred aspect, the AMV RT may comprise an α subunit reduced or substantially reduced in RNase H activity and a β subunit also reduced or substantially reduced in RNase H activity (*i.e.*, AMV α H⁻/ β H⁻ RT). In another preferred aspect, the AMV RT may comprise two β subunits, either or both of which may or may not be reduced or substantially reduced in RNase H activity (*i.e.*, AMV β H⁻/ β H⁻ RT; AMV β H⁻/ β H⁺ RT; or AMV β H⁺/ β H⁺ RT). In another preferred aspect, the AMV RT may comprise a single α subunit, which may or may not be reduced or substantially reduced in RNase H activity (*i.e.*, AMV α H⁻ RT or AMV α H⁺ RT). In another preferred aspect, the AMV RT may comprise one or more β p4 subunits, any or all of which may or may not be reduced or substantially reduced in RNase H activity (*e.g.*, AMV β p4H⁻/ β p4H⁻ RT; AMV

$\beta p4H/\beta p4H^+$ RT; AMV $\beta p4H^+/\beta p4H^+$ RT; AMV $\alpha H^-/\beta p4H^+$ RT; AMV $\alpha H^+/\beta p4H^+$ RT; AMV $\alpha H^-/\beta p4H^-$ RT; AMV $\alpha H^+/\beta p4H^-$ RT; AMV $\beta H^-/\beta p4H^-$ RT; AMV $\beta H^+/\beta p4H^-$ RT; etc.).

5 In a related aspect of the invention, where the AMV RTs comprise two or more subunits (or derivatives, variants, fragments or mutants thereof) and preferably comprise two subunits (*e.g.*, a dimer), at least one but preferably not all of these subunits may be modified or mutated to reduce, substantially reduce or eliminate the polymerase activity of at least one subunit (*e.g.*, pol-). In a preferred aspect, for an AMV RT which comprises an α and β subunit, the β subunit has been modified or mutated (preferably by recombinant techniques) to reduce, substantially reduce or eliminate the polymerase activity while the polymerase activity of the α subunit has not been mutated or modified in this manner. Preferably, the α subunit of such AMV RT has also been modified or mutated to reduce or substantially reduce RNase H activity while the β subunit has not been mutated or modified in this manner. Such a construct may be designated $\alpha H^-/\beta H^+$ pol-. Any number of combinations of subunits can be prepared in which mutations or modifications are made in one subunit of a two subunit enzyme, and these constructs may be combined with other modifications or mutations, such as those which reduce or substantially reduce RNase H activity. Illustrated examples include but are not limited to AMV $\alpha H^-/\beta H^-$ pol-; AMV $\alpha H^+/\beta H^+$ pol-; AMV αH^- pol-/ βH^+ ; AMV αH^- pol-/ βH^- ; AMV αH^+ pol-/ βH^+ ; AMV $\beta H^-/\beta H^-$ pol-; AMV $\beta H^-/\beta H^+$ pol-; AMV $\beta H^+/\beta H^+$ pol-; AMV $\beta p4H^-/\beta p4H^-$ pol-; AMV $\beta p4H^-/\beta p4H^+$ pol-; AMV $\beta p4H^+/\beta p4H^+$ pol-; AMV $\alpha H^-/\beta p4H^-$ pol-; AMV $\alpha H^+/\beta p4H^+$ pol-; AMV $\alpha H^-/\beta p4H^-$ pol-; AMV $\alpha H^+/\beta p4H^-$ pol-; AMV $\beta H^-/\beta p4H^+$ pol-; AMV $\beta H^-/\beta p4H^+$ pol-; AMV $\beta H^+/\beta p4H^-$ pol-; etc. In a preferred aspect, the polymerase domain of the subunit is modified or mutated (one or more point mutations, deletion mutations and/or insertion mutations) by recombinant techniques. In another aspect, the nucleotide binding site of the polymerase domain is modified or mutated. In a preferred aspect, one or more acidic amino

10
15
20
25

acids within the nucleotide binding site are substituted with different amino acids. Particularly preferred amino acids within the nucleotide binding site for mutation or modification include, but are not limited to, Asp¹⁰⁷, Leu¹⁰⁸, Lys¹⁰⁹, and Asp¹¹⁰, or the corresponding amino acid sequence.

5 The invention also relates to methods of producing the AMV RTs of the invention, which methods comprise obtaining a host cell comprising a nucleic acid sequence encoding one or more α subunits (or derivatives, variants, fragments or mutants thereof) and/or a nucleic acid sequence encoding one or more β subunits (or derivatives, variants, fragments or mutants thereof) and/or a nucleic acid sequence encoding one or more β p4 subunits (or derivatives, variants, fragments or mutants thereof), and culturing the host cell under conditions sufficient to produce the AMV RTs. The nucleic acid sequences encoding such α subunit(s) and/or such β subunit(s) and/or such β p4 subunit(s) may be contained in the same vector or in different vectors. In accordance with the invention, such α subunit(s) and/or β subunit(s) and/or β p4 subunit(s) may be produced separately and mixed before or after isolation of each subunit to form the AMV RTs of the invention. Alternatively, such α and/or β and/or β p4 subunits may be expressed simultaneously (*i.e.*, co-expressed) in the same host cell, thereby producing an AMV RT comprising an α and a β subunit, an α subunit alone, a β subunit alone, a β p4 subunit alone, a β subunit and a β p4 subunit, two β subunits, or two β p4 subunits, or derivatives, variants, fragments or mutants thereof. In a preferred aspect, the α subunit is simultaneously expressed in a host cell with the β subunit. In a related aspect of the invention, the β or β p4 subunits (or derivatives, variants, fragments or mutants thereof) may be expressed in a host or host cell which accomplishes *in vivo* processing as described above for production of RSV RTs. Thus, by expressing a β subunit and/or a β p4 subunit (or derivatives, fragments or mutants thereof) in a host system which has appropriate processing enzymes or proteins, the corresponding α subunit may be produced allowing formation of an AMV RT which comprises an α and a β subunit, an α and a β p4 subunit, etc.

As noted above, any or all of the α and/or β and/or $\beta p4$ subunits (or derivatives, variants, fragments or mutants thereof) of the AMV RT may be reduced or substantially reduced in RNase H activity. These AMV RTs or subunits thereof may then be isolated from the host cell and may be substantially purified by those methods described above for purification of RSV RTs. The invention also relates to kits comprising one or more of the AMV RTs of the invention.

The invention also relates to other substantially pure ASLV reverse transcriptases, which may or may not be reduced or substantially reduced in RNase H activity. Such ASLV RTs may comprise one or more subunits (or derivatives, variants, fragments or mutants thereof) selected from one or more α subunits, one or more β subunits, and one or more $\beta p4$ subunits, as described for RSV RT and AMV RT above. The invention also relates to methods of producing such ASLV RTs of the invention as described above for RSV RT and AMV RT.

The invention also concerns the RSV reverse transcriptases and AMV reverse transcriptases or other ASLV reverse transcriptases and subunits thereof (and derivatives, variants, fragments and mutants thereof) of the invention which have functional activity, as measured by the ability of the proteins to produce first strand cDNA from a mRNA template. Such functional activity may be measured in accordance with the invention based on the total full-length reverse transcribed product made during the synthesis reaction. The amount of product is preferably measured based on the mass (*e.g.*, nanograms) of products produced, although other means of measuring the amount of product will be recognized by one of ordinary skill in the art. Additionally, functional activity may be measured in terms of the percentage of full-length products produced during a cDNA synthesis reaction. For example, the percent full-length functional activity may be determined by dividing the amount of full-length product by the amount of total product produced during a cDNA synthesis reaction and multiplying the result by 100 to obtain the percentage. The RSV and AMV reverse transcriptases and their subunits (and derivatives, variants, fragments and mutants thereof) of the invention

produce greater than about 4%, preferably greater than about 5%, more preferably greater than about 7.5%, still more preferably greater than about 10%, still more preferably greater than about 20%, and most preferably greater than about 25%, full-length cDNA in a nucleic acid synthesis reaction. Preferred ranges of such percentages include about 5% to about 100%, about 7.5% to about 75%, about 7.5% to about 50%, about 10% to about 50%, about 15% to about 40%, about 20% to about 40%, and about 20% to about 50%. Functional activity may also be measured in accordance with the invention by determining the percentage of total cDNA product compared to the amount of input mRNA in the synthesis reaction. Thus, the total amount of cDNA product is divided by the amount of input mRNA, the result of which is multiplied by 100 to determine the percentage functional activity associated with amount of product produced compared to amount of the template used. Preferably, the reverse transcriptases of the invention produce greater than about 15%, more preferably greater than about 20%, still more preferably greater than about 25%, still more preferably greater than about 30%, and most preferably greater than about 40%, of cDNA compared to input mRNA in the cDNA synthesis reaction. Preferred ranges of such percentages include about 5% to about 100%, about 10% to about 80%, about 15% to about 80%, about 15% to about 75%, about 20% to about 75%, about 20% to about 70%, about 25% to about 75%, about 25% to about 70%, about 25% to about 60%, and about 25% to about 50%. The AMV and RSV reverse transcriptases and their subunits (and derivatives, variants, fragments and mutants thereof) of the invention preferably have specific activities greater than about 5 units/mg, more preferably greater than about 50 units/mg, still more preferably greater than about 100 units/mg, 250 units/mg, 500 units/mg, 1000 units/mg, 5000 units/mg or 10,000 units/mg, and most preferably greater than about 15,000 units/mg, greater than about 16,000 units/mg, greater than about 17,000 units/mg, greater than about 18,000 units/mg, greater than about 19,000 units/mg and greater than about 20,000 units/mg. Preferred ranges of specific activities for

the AMV and RSV RTs and their subunits (or derivatives, variants, fragments or mutants thereof) of the invention include a specific activity from about 5 units/mg to about 140,000 units/mg, a specific activity from about 5 units/mg to about 125,000 units/mg, a specific activity of from about 50 units/mg to about 100,000 units/mg, a specific activity from about 100 units/mg to about 100,000 units/mg, a specific activity from about 250 units/mg to about 100,000 units/mg, a specific activity from about 500 units/mg to about 100,000 units/mg, a specific activity from about 1000 units/mg to about 100,000 units/mg, a specific activity from about 5000 units/mg to about 100,000 units/mg, a specific activity from about 10,000 units/mg to about 100,000 units/mg, a specific activity from about 25,000 units/mg to about 75,000 units/mg. Other preferred ranges of specific activities include a specific activity of from about 20,000 units/mg to about 140,000 units/mg, a specific activity from about 20,000 units/mg to about 130,000 units/mg, a specific activity from about 20,000 units/mg to about 120,000 units/mg, a specific activity from about 20,000 units/mg to about 110,000 units/mg, a specific activity from about 20,000 units/mg to about 100,000 units/mg, a specific activity from about 20,000 units/mg to about 90,000 units/mg, a specific activity from about 25,000 units/mg to about 140,000 units/mg, a specific activity from about 25,000 units/mg to about 130,000 units/mg, a specific activity from about 25,000 units/mg to about 120,000 units/mg, a specific activity from about 25,000 units/mg to about 110,000 units/mg, a specific activity from about 25,000 units/mg to about 100,000 units/mg, and a specific activity from about 25,000 units/mg to about 90,000 units/mg. Preferably, the lower end of the specific activity range may vary from 30,000, 35,000, 40,000, 45,000, 50,000, 5,000, 60,000, 65,000, 70,000, 75,000, and 80,000 units/mg, while the upper end of the range may vary from 150,000, 140,000, 130,000, 120,000, 110,000, 100,000, and 90,000 units/mg. In accordance with the invention, specific activity is a measurement of the enzymatic activity (in units) of the protein or enzyme relative to the total amount of protein or enzyme used in a reaction. The

measurement of a specific activity may be determined by standard techniques well-known to one of ordinary skill in the art. Preferred assays for determining the specific activity of an enzyme or protein are described in detail in the Examples below.

5 The RSV RTs and AMV RTs or other ASLV RTs and their subunits (or derivatives, variants, fragments or mutants thereof) of the invention may be used to make nucleic acid molecules from one or more templates. Such methods comprise mixing one or more nucleic acid templates (*e.g.*, mRNA, and more preferably a population of mRNA molecules) with one or more of the RSV RTs and/or one or more AMV RTs and/or other ASLV RTs of the invention and
10 incubating the mixture under conditions sufficient to make one or more nucleic acid molecules complementary to all or a portion of the one or more nucleic acid templates.

 The invention also relates to methods for the amplification of one or more nucleic acid molecules comprising mixing one or more nucleic acid templates with
15 one or more of the RSV RTs and/or one or more of the AMV RTs and/or other ASLV RTs of the invention and optionally with one or more DNA polymerases, and incubating the mixture under conditions sufficient to amplify one or more nucleic acid molecules complementary to all or a portion of the one or more
20 nucleic acid templates.

 The invention also concerns methods for the sequencing of one or more nucleic acid molecules comprising (a) mixing one or more nucleic acid molecules to be sequenced with one or more primer nucleic acid molecules, one or more RSV RTs and/or one or more AMV RTs and/or other ASLV RTs of the
25 invention, one or more nucleotides and one or more terminating agents; (b) incubating the mixture under conditions sufficient to synthesize a population of nucleic acid molecules complementary to all or a portion of the one or more nucleic acid molecules to be sequenced; and (c) separating the population of

nucleic acid molecules to determine the nucleotide sequence of all or a portion of the one or more nucleic acid molecules to be sequenced.

The invention also concerns methods for elevated- or high-temperature reverse transcription of a nucleic acid molecule comprising (a) mixing a nucleic acid template (preferably an RNA (*e.g.*, a mRNA molecule or a polyA⁺ RNA molecule) or a DNA molecule) with one or more polypeptides having reverse transcriptase activity; and (b) incubating the mixture at a temperature of 50°C or greater and under conditions sufficient to make a first nucleic acid molecule (such as a full length cDNA molecule) complementary to all or a portion of the nucleic acid template. In preferred such methods, the mixture is incubated at a temperature of about 51°C or greater, about 52°C or greater, about 53°C or greater, about 54°C or greater, about 55°C or greater, about 56°C or greater, about 57°C or greater, about 58°C or greater, about 59°C or greater, about 60°C or greater, about 61°C or greater, about 62°C or greater; or at a temperature range of from about 50°C to about 70°C, about 51°C to about 70°C, about 52°C to about 70°C, about 53°C to about 70°C, about 54°C to about 70°C, about 55°C to about 70°C, about 55°C to about 65°C, about 56°C to about 65°C, about 56°C to about 64°C or about 56°C to about 62°C. The invention is also directed to such methods which further comprise incubating the first nucleic acid molecule under conditions sufficient to make a second nucleic acid molecule complementary to all or portion of the first nucleic acid molecule. According to the invention, the first and second nucleic acid molecules produced by these methods may be DNA molecules, and may form a double stranded DNA molecule which may be a full length cDNA molecule. The one or more polypeptides having reverse transcriptase activity that are used in these methods preferably are reduced or substantially reduced in RNase H activity, and may be selected from the group consisting of one or more subunits of AMV reverse transcriptase and one or more subunits of RSV reverse transcriptase and one or more subunits of other ASLV reverse transcriptases (or derivatives, variants, fragments or mutants thereof).

As noted above, such AMV RTs or RSV RTs or other ASLV RTs may comprise one or more α subunits, one or more β subunits, and/or one or more β p4 subunits, any or all of which subunits may be reduced or substantially reduced in RNase H activity. Particularly preferred polymerases having RT activity for use in these methods are those RSV RTs and AMV RTs or other ASLV RTs and their subunits (or derivatives, variants, fragments or mutants thereof) described above.

The invention also concerns nucleic acid molecules produced by such methods (which may be full-length cDNA molecules), vectors (particularly expression vectors) comprising these nucleic acid molecules and host cells comprising these vectors and nucleic acid molecules.

Sources of Enzymes

Enzymes for use in the compositions, methods and kits of the invention include any enzyme having reverse transcriptase activity. Such enzymes include, but are not limited to, retroviral reverse transcriptase, retrotransposon reverse transcriptase, hepatitis B reverse transcriptase, cauliflower mosaic virus reverse transcriptase, bacterial reverse transcriptase, *Tth* DNA polymerase, *Taq* DNA polymerase (Saiki, R.K., *et al.*, *Science* 239:487-491 (1988); U.S. Patent Nos. 4,889,818 and 4,965,188), *Tne* DNA polymerase (WO 96/10640), *Tma* DNA polymerase (U. S. Patent No. 5,374,553) and mutants, fragments, variants or derivatives thereof (*see, e.g.*, commonly owned, co-pending U.S. Patent Application Nos. 08/706,702 and 08/706,706, both filed September 9, 1996, which are incorporated by reference herein in their entireties). As will be understood by one of ordinary skill in the art, modified reverse transcriptases may be obtained by recombinant or genetic engineering techniques that are routine and well-known in the art. Mutant reverse transcriptases can, for example, be obtained by mutating the gene or genes encoding the reverse transcriptase of interest by site-directed or random mutagenesis. Such mutations may include point mutations, deletion mutations and insertional mutations. Preferably, one or

more point mutations (*e.g.*, substitution of one or more amino acids with one or more different amino acids) are used to construct mutant reverse transcriptases of the invention. Fragments of reverse transcriptases may be obtained by deletion mutation by recombinant techniques that are routine and well-known in the art, or
5 by enzymatic digestion of the reverse transcriptase(s) of interest using any of a number of well-known proteolytic enzymes.

Preferred enzymes for use in the invention include those that are reduced or substantially reduced in RNase H activity. Such enzymes that are reduced or substantially reduced in RNase H activity may be obtained by mutating the RNase
10 H domain within the reverse transcriptase of interest, preferably by one or more point mutations, one or more deletion mutations, and/or one or more insertion mutations as described above. By an enzyme "substantially reduced in RNase H activity" is meant that the enzyme has less than about 30%, less than about 25%,
15 20%, more preferably less than about 15%, less than about 10%, less than about 7.5%, or less than about 5%, and most preferably less than about 5% or less than about 2%, of the RNase H activity of the corresponding wildtype or RNase H⁺ enzyme such as wildtype Moloney Murine Leukemia Virus (M-MLV), Avian Myeloblastosis Virus (AMV) or Rous Sarcoma Virus (RSV) reverse transcriptases. The RNase H activity of any enzyme may be determined by a
20 variety of assays, such as those described, for example, in U.S. Patent No. 5,244,797, in Kotewicz, M.L., *et al.*, *Nucl. Acids Res.* 16:265 (1988), in Gerard, G.F., *et al.*, *FOCUS* 14(5):91 (1992), and in U.S. Patent No. 5,668,005, the disclosures of all of which are fully incorporated herein by reference.

Particularly preferred enzymes for use in the invention include, but are not
25 limited to, M-MLV H⁻ reverse transcriptase, RSV H⁻ reverse transcriptase, AMV H⁻ reverse transcriptase, RAV H⁻ reverse transcriptase, MAV H⁻ reverse transcriptase and HIV H⁻ reverse transcriptase. It will be understood by one of ordinary skill, however, that any enzyme capable of producing a DNA molecule from a ribonucleic acid molecule (*i.e.*, having reverse transcriptase activity) that

is substantially reduced in RNase H activity may be equivalently used in the compositions, methods and kits of the invention.

Enzymes used in the invention may have distinct reverse transcription pause sites with respect to the template nucleic acid. Whether or not two enzymes have distinct reverse transcription pause sites may be determined by a variety of assays, including, for example, electrophoretic analysis of the chain lengths of DNA molecules produced by the two enzymes (Weaver, D.T., and DePamphilis, M.L., *J. Biol. Chem.* 257(4):2075-2086 (1982); Abbots, J., *et al.*, *J. Biol. Chem.* 268(14):10312-10323 (1993)), or by other assays that will be familiar to one of ordinary skill in the art. As described above, these distinct transcription pause sites may represent secondary structural and sequence barriers in the nucleic acid template which occur frequently at homopolymer stretches. Thus, for example, the second enzyme may reverse transcribe to a point (*e.g.*, a hairpin) on the template nucleic acid that is proximal or distal (*i.e.*, 3' or 5') to the point to which the first enzyme reverse transcribes the template nucleic acid. This combination of two or more enzymes having distinct reverse transcription pause sites facilitates production of full-length cDNA molecules since the secondary structural and sequence barriers may be overcome. Moreover, the elevated- or high-temperature reverse transcription of the invention may also assist in overcoming secondary structural and sequence barriers during nucleic acid synthesis. Thus, the elevated- or high-temperature synthesis may be used in combination with the two or more reverse transcriptases (preferably using an AMV RT, an RSV RT or other ASLV RT) to facilitate full-length cDNA synthesis.

A variety of DNA polymerases are useful in accordance with the present invention. Such polymerases include, but are not limited to, *Thermus thermophilus* (*Tth*) DNA polymerase, *Thermus aquaticus* (*Taq*) DNA polymerase, *Thermotoga neapolitana* (*Tne*) DNA polymerase, *Thermotoga maritima* (*Tma*) DNA polymerase, *Thermococcus litoralis* (*Tli* or VENT™) DNA polymerase, *Pyrococcus furiosus* (*Pfu*) DNA polymerase, DEEPVENT™ DNA polymerase,

Pyrococcus woosii (*Pwo*) DNA polymerase, *Bacillus sterothermophilus* (*Bst*) DNA polymerase, *Bacillus caldophilus* (*Bca*) DNA polymerase, *Sulfolobus acidocaldarius* (*Sac*) DNA polymerase, *Thermoplasma acidophilum* (*Tac*) DNA polymerase, *Thermus flavus* (*Tfl/Tub*) DNA polymerase, *Thermus ruber* (*Tru*) DNA polymerase, *Thermus brockianus* (DYNAZYME™) DNA polymerase, *Methanobacterium thermoautotrophicum* (*Mth*) DNA polymerase, *Mycobacterium* spp. DNA polymerase (*Mtb*, *Mlep*), and mutants, variants and derivatives thereof.

DNA polymerases used in accordance with the invention may be any enzyme that can synthesize a DNA molecule from a nucleic acid template, typically in the 5' to 3' direction. Such polymerases may be mesophilic or thermophilic, but are preferably thermophilic. Mesophilic polymerases include T5 DNA polymerase, T7 DNA polymerase, Klenow fragment DNA polymerase, DNA polymerase III, and the like. Preferred DNA polymerases are thermostable DNA polymerases such as *Taq*, *Tne*, *Tma*, *Pfu*, VENT™, DEEPVENT™, *Tth* and mutants, variants and derivatives thereof (U.S. Patent No. 5,436,149; U.S. Patent No. 5,512,462; WO 92/06188; WO 92/06200; WO 96/10640; Barnes, W.M., *Gene* 112:29-35 (1992); Lawyer, F.C., *et al.*, *PCR Meth. Appl.* 2:275-287 (1993); Flaman, J.-M., *et al.*, *Nucl. Acids Res.* 22(15):3259-3260 (1994)). For amplification of long nucleic acid molecules (*e.g.*, nucleic acid molecules longer than about 3-5 Kb in length), at least two DNA polymerases (one substantially lacking 3' exonuclease activity and the other having 3' exonuclease activity) are typically used. See U.S. Patent No. 5,436,149; U.S. Patent No. 5,512,462; Barnes, W.M., *Gene* 112:29-35 (1992); and commonly owned, co-pending U.S. Patent Application No. 08/801,720, filed February 14, 1997, the disclosures of all of which are incorporated herein in their entireties. Examples of DNA polymerases substantially lacking in 3' exonuclease activity include, but are not limited to, *Taq*, *Tne*(exo⁻), *Tma*, *Pfu*(exo⁻), *Pwo* and *Tth* DNA polymerases, and mutants, variants and derivatives thereof. Nonlimiting examples of DNA

polymerases having 3' exonuclease activity include *Pfu*/DEEPVENT™ and *Tli*/VENT™ and mutants, variants and derivatives thereof.

Polypeptides having reverse transcriptase activity for use in the invention may be obtained commercially, for example from Life Technologies, Inc. (Rockville, Maryland), Pharmacia (Piscataway, New Jersey), Sigma (Saint Louis, Missouri) or Boehringer Mannheim Biochemicals (Indianapolis, Indiana). Alternatively, polypeptides having reverse transcriptase activity may be isolated from their natural viral or bacterial sources according to standard procedures for isolating and purifying natural proteins that are well-known to one of ordinary skill in the art (*see, e.g.,* Houts, G.E., *et al., J. Virol.* 29:517 (1979)). In addition, the polypeptides having reverse transcriptase activity may be prepared by recombinant DNA techniques that are familiar to one of ordinary skill in the art (*see, e.g.,* Kotewicz, M.L., *et al., Nucl. Acids Res.* 16:265 (1988); Soltis, D.A., and Skalka, A.M., *Proc. Natl. Acad. Sci. USA* 85:3372-3376 (1988)).

DNA polymerases for use in the invention may be obtained commercially, for example from Life Technologies, Inc. (Rockville, Maryland), Perkin-Elmer (Branchburg, New Jersey), New England BioLabs (Beverly, Massachusetts) or Boehringer Mannheim Biochemicals (Indianapolis, Indiana).

Formulation of Enzyme Compositions

To form the compositions of the present invention, two or more reverse transcriptases are preferably admixed in a buffered salt solution. One or more DNA polymerases and/or one or more nucleotides may optionally be added to make the compositions of the invention. More preferably, the enzymes are provided at working concentrations in stable buffered salt solutions. The terms "stable" and "stability" as used herein generally mean the retention by a composition, such as an enzyme composition, of at least 70%, preferably at least 80%, and most preferably at least 90%, of the original enzymatic activity (in units) after the enzyme or composition containing the enzyme has been stored for about

one week at a temperature of about 4°C, about two to six months at a temperature of about -20°C, and about six months or longer at a temperature of about -80°C. As used herein, the term "working concentration" means the concentration of an enzyme that is at or near the optimal concentration used in a solution to perform a particular function (such as reverse transcription of nucleic acids).

The water used in forming the compositions of the present invention is preferably distilled, deionized and sterile filtered (through a 0.1-0.2 micrometer filter), and is free of contamination by DNase and RNase enzymes. Such water is available commercially, for example from Sigma Chemical Company (Saint Louis, Missouri), or may be made as needed according to methods well known to those skilled in the art.

In addition to the enzyme components, the present compositions preferably comprise one or more buffers and cofactors necessary for synthesis of a nucleic acid molecule such as a cDNA molecule. Particularly preferred buffers for use in forming the present compositions are the acetate, sulfate, hydrochloride, phosphate or free acid forms of Tris-(hydroxymethyl)aminomethane (TRIS®), although alternative buffers of the same approximate ionic strength and pKa as TRIS® may be used with equivalent results. In addition to the buffer salts, cofactor salts such as those of potassium (preferably potassium chloride or potassium acetate) and magnesium (preferably magnesium chloride or magnesium acetate) are included in the compositions. Addition of one or more carbohydrates and/or sugars to the compositions and/or synthesis reaction mixtures may also be advantageous, to support enhanced stability of the compositions and/or reaction mixtures upon storage. Preferred such carbohydrates or sugars for inclusion in the compositions and/or synthesis reaction mixtures of the invention include, but are not limited to, sucrose, trehalose, and the like. Furthermore, such carbohydrates and/or sugars may be added to the storage buffers for the enzymes used in the production of the enzyme compositions and kits of the invention. Such

carbohydrates and/or sugars are commercially available from a number of sources, including Sigma (St. Louis, MO).

5 It is often preferable to first dissolve the buffer salts, cofactor salts and carbohydrates or sugars at working concentrations in water and to adjust the pH of the solution prior to addition of the enzymes. In this way, the pH-sensitive enzymes will be less subject to acid- or alkaline-mediated inactivation during formulation of the present compositions.

10 To formulate the buffered salts solution, a buffer salt which is preferably a salt of Tris(hydroxymethyl)aminomethane (TRIS®), and most preferably the hydrochloride salt thereof, is combined with a sufficient quantity of water to yield a solution having a TRIS® concentration of 5-150 millimolar, preferably 10-60 millimolar, and most preferably about 20-60 millimolar. To this solution, a salt of magnesium (preferably either the chloride or acetate salt thereof) may be added to provide a working concentration thereof of 1-10 millimolar, preferably 1.5-8.0 millimolar, and most preferably about 3-7.5 millimolar. A salt of potassium (preferably a chloride or acetate salt of potassium) may also be added to the solution, at a working concentration of 10-100 millimolar and most preferably about 75 millimolar. A reducing agent such as dithiothreitol may be added to the solution, preferably at a final concentration of about 1-100 mM, more preferably a concentration of about 5-50 mM or about 7.5-20 mM, and most preferably at a concentration of about 10 mM. Preferred concentrations of carbohydrates and/or sugars for inclusion in the compositions of the invention range from about 5% (w/v) to about 30% (w/v), about 7.5% (w/v) to about 25% (w/v), about 10% (w/v) to about 25% (w/v), about 10% (w/v) to about 20% (w/v), and preferably about 10% (w/v) to about 15% (w/v). A small amount of a salt of ethylenediaminetetraacetate (EDTA), such as disodium EDTA, may also be added (preferably about 0.1 millimolar), although inclusion of EDTA does not appear to be essential to the function or stability of the compositions of the present invention. After addition of all buffers and salts, this buffered salt solution is

15

20

25

mixed well until all salts are dissolved, and the pH is adjusted using methods known in the art to a pH value of 7.4 to 9.2, preferably 8.0 to 9.0, and most preferably about 8.4.

To these buffered salt solutions, the enzymes (reverse transcriptases and/or DNA polymerases) are added to produce the compositions of the present invention. M-MLV RTs are preferably added at a working concentration in the solution of about 1,000 to about 50,000 units per milliliter, about 2,000 to about 30,000 units per milliliter, about 2,500 to about 25,000 units per milliliter, about 3,000 to about 22,500 units per milliliter, about 4,000 to about 20,000 units per milliliter, and most preferably at a working concentration of about 5,000 to about 20,000 units per milliliter. AMV RTs, MAV RTs, RSV RTs and RAV RTs, including those of the invention described above, are preferably added at a working concentration in the solution of about 100 to about 5000 units per milliliter, about 125 to about 4000 units per milliliter, about 150 to about 3000 units per milliliter, about 200 to about 2500 units per milliliter, about 225 to about 2000 units per milliliter, and most preferably at a working concentration of about 250 to about 1000 units per milliliter. The enzymes in the thermophilic DNA polymerase group (*Taq*, *Tne*, *Tma*, *Pfu*, VENT™, DEEPVENT™, *Tth* and mutants, variants and derivatives thereof) are preferably added at a working concentration in the solution of about 100 to about 1000 units per milliliter, about 125 to about 750 units per milliliter, about 150 to about 700 units per milliliter, about 200 to about 650 units per milliliter, about 225 to about 550 units per milliliter, and most preferably at a working concentration of about 250 to about 500 units per milliliter. The enzymes may be added to the solution in any order, or may be added simultaneously.

The compositions of the invention may further comprise one or more nucleotides, which are preferably deoxynucleoside triphosphates (dNTPs) or dideoxynucleoside triphosphates (ddNTPs). The dNTP components of the present compositions serve as the "building blocks" for newly synthesized nucleic acids,

being incorporated therein by the action of the polymerases, and the ddNTPs may be used in sequencing methods according to the invention. Examples of nucleotides suitable for use in the present compositions include, but are not limited to, dUTP, dATP, dTTP, dCTP, dGTP, dITP, 7-deaza-dGTP, α -thio-dATP, α -thio-dTTP, α -thio-dGTP, α -thio-dCTP, ddUTP, ddATP, ddTTP, ddCTP, ddGTP, ddITP, 7-deaza-ddGTP, α -thio-ddATP, α -thio-ddTTP, α -thio-ddGTP, α -thio-ddCTP or derivatives thereof, all of which are available commercially from sources including Life Technologies, Inc. (Rockville, Maryland), New England BioLabs (Beverly, Massachusetts) and Sigma Chemical Company (Saint Louis, Missouri). The nucleotides may be unlabeled, or they may be detectably labeled by coupling them by methods known in the art with radioisotopes (*e.g.*, ^3H , ^{14}C , ^{32}P or ^{35}S), vitamins (*e.g.*, biotin), fluorescent moieties (*e.g.*, fluorescein, rhodamine, Texas Red, or phycoerythrin), chemiluminescent labels (*e.g.*, using the PHOTO-GENETM or ACESTM chemiluminescence systems, available commercially from Life Technologies, Inc., Rockville, Maryland), dioxigenin and the like. Labeled nucleotides may also be obtained commercially, for example from Life Technologies, Inc. (Rockville, Maryland) or Sigma Chemical Company (Saint Louis, Missouri). In the present compositions, the nucleotides are added to give a working concentration of each nucleotide of about 10-4000 micromolar, about 50-2000 micromolar, about 100-1500 micromolar, or about 200-1200 micromolar, and most preferably a concentration of about 1000 micromolar.

To reduce component deterioration, storage of the reagent compositions is preferably at about 4°C for up to one day, or most preferably at -20°C for up to one year.

In another aspect, the compositions and reverse transcriptases of the invention may be prepared and stored in dry form in the presence of one or more carbohydrates, sugars, or synthetic polymers. Preferred carbohydrates, sugars or polymers for the preparation of dried compositions or reverse transcriptases include, but are not limited to, sucrose, trehalose, and polyvinylpyrrolidone (PVP)

or combinations thereof. *See, e.g.*, U.S. Patent Nos. 5,098,893, 4,891,319, and 5,556,771, the disclosures of which are entirely incorporated herein by reference. Such dried compositions and enzymes may be stored at various temperatures for extended times without significant deterioration of enzymes or components of the compositions of the invention. Preferably, the dried reverse transcriptases or compositions are stored at 4°C or at -20°C.

Production of cDNA Molecules

Sources of Nucleic Acid Molecules

In accordance with the invention, cDNA molecules (single-stranded or double-stranded) may be prepared from a variety of nucleic acid template molecules. Preferred nucleic acid molecules for use in the present invention include single-stranded or double-stranded DNA and RNA molecules, as well as double-stranded DNA:RNA hybrids. More preferred nucleic acid molecules include messenger RNA (mRNA), transfer RNA (tRNA) and ribosomal RNA (rRNA) molecules, although mRNA molecules are the preferred template according to the invention.

The nucleic acid molecules that are used to prepare cDNA molecules according to the methods of the present invention may be prepared synthetically according to standard organic chemical synthesis methods that will be familiar to one of ordinary skill. More preferably, the nucleic acid molecules may be obtained from natural sources, such as a variety of cells, tissues, organs or organisms. Cells that may be used as sources of nucleic acid molecules may be prokaryotic (bacterial cells, including but not limited to those of species of the genera *Escherichia*, *Bacillus*, *Serratia*, *Salmonella*, *Staphylococcus*, *Streptococcus*, *Clostridium*, *Chlamydia*, *Neisseria*, *Treponema*, *Mycoplasma*, *Borrelia*, *Legionella*, *Pseudomonas*, *Mycobacterium*, *Helicobacter*, *Erwinia*, *Agrobacterium*, *Rhizobium*, *Xanthomonas* and *Streptomyces*) or eukaryotic (including fungi (especially yeasts), plants, protozoans and other parasites, and

animals including insects (particularly *Drosophila* spp. cells), nematodes (particularly *Caenorhabditis elegans* cells), and mammals (particularly human cells)).

5 Mammalian somatic cells that may be used as sources of nucleic acids include blood cells (reticulocytes and leukocytes), endothelial cells, epithelial cells, neuronal cells (from the central or peripheral nervous systems), muscle cells (including myocytes and myoblasts from skeletal, smooth or cardiac muscle), connective tissue cells (including fibroblasts, adipocytes, chondrocytes, chondroblasts, osteocytes and osteoblasts) and other stromal cells (e.g.,
10 macrophages, dendritic cells, Schwann cells). Mammalian germ cells (spermatocytes and oocytes) may also be used as sources of nucleic acids for use in the invention, as may the progenitors, precursors and stem cells that give rise to the above somatic and germ cells. Also suitable for use as nucleic acid sources are mammalian tissues or organs such as those derived from brain, kidney, liver,
15 pancreas, blood, bone marrow, muscle, nervous, skin, genitourinary, circulatory, lymphoid, gastrointestinal and connective tissue sources, as well as those derived from a mammalian (including human) embryo or fetus.

Any of the above prokaryotic or eukaryotic cells, tissues and organs may be normal, diseased, transformed, established, progenitors, precursors, fetal or
20 embryonic. Diseased cells may, for example, include those involved in infectious diseases (caused by bacteria, fungi or yeast, viruses (including AIDS, HIV, HTLV, herpes, hepatitis and the like) or parasites), in genetic or biochemical pathologies (e.g., cystic fibrosis, hemophilia, Alzheimer's disease, muscular dystrophy or multiple sclerosis) or in cancerous processes. Transformed or established animal
25 cell lines may include, for example, COS cells, CHO cells, VERO cells, BHK cells, HeLa cells, HepG2 cells, K562 cells, 293 cells, L929 cells, F9 cells, and the like. Other cells, cell lines, tissues, organs and organisms suitable as sources of nucleic acids for use in the present invention will be apparent to one of ordinary skill in the art.

Once the starting cells, tissues, organs or other samples are obtained, nucleic acid molecules (such as mRNA) may be isolated therefrom by methods that are well-known in the art (*See, e.g.,* Maniatis, T., *et al.*, *Cell* 15:687-701 (1978); Okayama, H., and Berg, P., *Mol. Cell. Biol.* 2:161-170 (1982); Gubler, U., and Hoffman, B.J., *Gene* 25:263-269 (1983)). The nucleic acid molecules thus isolated may then be used to prepare cDNA molecules and cDNA libraries in accordance with the present invention.

In the practice of the invention, cDNA molecules or cDNA libraries are produced by mixing one or more nucleic acid molecules obtained as described above, which is preferably one or more mRNA molecules such as a population of mRNA molecules, with two or more polypeptides having reverse transcriptase activity, or with one or more of the compositions of the invention or with one or more of the RSV RTs and/or AMV RTs and/or other ASLV RTs of the invention, under conditions favoring the reverse transcription of the nucleic acid molecule by the action of the enzymes or the compositions to form a cDNA molecule (single-stranded or double-stranded). Thus, the method of the invention comprises (a) mixing one or more nucleic acid templates (preferably one or more RNA or mRNA templates, such as a population of mRNA molecules) with one or more reverse transcriptases of the invention and (b) incubating the mixture under conditions sufficient to make one or more nucleic acid molecules complementary to all or a portion of the one or more templates. Such methods may include the use of one or more DNA polymerases. The invention may be used in conjunction with methods of cDNA synthesis such as those described in the Examples below, or others that are well-known in the art (*see, e.g.,* Gubler, U., and Hoffman, B.J., *Gene* 25:263-269 (1983); Krug, M.S., and Berger, S.L., *Meth. Enzymol.* 152:316-325 (1987); Sambrook, J., *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, pp. 8.60-8.63 (1989)), to produce cDNA molecules or libraries.

The invention is also particularly directed to methods for reverse transcription of a nucleic acid molecule at elevated temperatures. As described in more detail in Example 5, retroviral RTs are generally not used at temperatures above 37°C to 42°C to copy nucleic acid templates such as RNA molecules because of the limited thermal stability of these mesophilic enzymes. At these temperatures, however, mRNA secondary structure may interfere with reverse transcription (Gerard, G.F., *et al.*, *FOCUS* 11:60 (1989); Myers, T.W., and Gelfand, D.H., *Biochem.* 30:7661 (1991)), and the specificity of primer binding may be reduced during gene-specific reverse transcription processes, such as RT-PCR, causing high background signal (Myers, T.W., and Gelfand, D.H., *Biochem.* 30:7661 (1991); Freeman, W.N., *et al.*, *BioTechniques* 20:782 (1996)). To help overcome these problems, the present invention therefore provides methods of RNA reverse transcription at more elevated temperatures, *i.e.*, above 50°C.

Therefore, the invention is related to methods for reverse transcription of a nucleic acid molecule comprising (a) mixing a nucleic acid template with one or more polypeptides having reverse transcriptase activity; and (b) incubating the mixture at a temperature of about 50°C or greater and under conditions sufficient to make a first nucleic acid molecule complementary to all or a portion of the nucleic acid template. Nucleic acid templates which may be copied according to these methods include, but are not limited to, an RNA molecule (*e.g.*, a mRNA molecule or a polyA⁺ RNA molecule) and a DNA molecule (*e.g.*, a single-stranded or double-stranded DNA molecule). According to the invention, the first nucleic acid molecule produced by these methods may be a full length cDNA molecule. While any incubation temperature of about 50°C or greater may be used in the present methods, particularly preferred incubation temperatures include, but are not limited to, temperatures of about 51°C or greater, about 52°C or greater, about 53°C or greater, about 54°C or greater, about 55°C or greater, about 56°C or greater, about 57°C or greater, about 58°C or greater, about 59°C or greater, about 60°C or greater, about 61°C or greater, about 62°C or greater,

about 63°C or greater, about 64°C or greater, about 65°C or greater, about 66°C or greater, about 67°C or greater, about 68°C or greater, about 69°C or greater or about 70°C or greater. In other such methods, the incubation temperature may be over a range of incubation temperatures, including but not limited to a temperature range of from about 50°C to about 70°C, about 51°C to about 70°C, about 52°C to about 70°C, about 53°C to about 70°C, about 54°C to about 70°C, about 55°C to about 70°C, about 55°C to about 69°C, about 55°C to about 68°C, about 55°C to about 67°C, about 55°C to about 66°C, about 55°C to about 65°C, about 56°C to about 65°C, about 56°C to about 64°C or about 56°C to about 62°C. The invention is also directed to such methods which further comprise incubating the first nucleic acid molecule under conditions sufficient to make a second nucleic acid molecule complementary to all or portion of the first nucleic acid molecule. According to the invention, the first and second nucleic acid molecules produced by these methods may be DNA molecules, and may form a double stranded DNA molecule which may be a full length cDNA molecule. As described for the methods above, the one or more polypeptides having reverse transcriptase activity that are used in these higher-temperature methods preferably are reduced or substantially reduced in RNase H activity, and may be selected from the group consisting of one or more AMV reverse transcriptases or subunits thereof (or derivatives, variants, fragments or mutants thereof), and one or more RSV reverse transcriptases or subunits thereof (or derivatives, variants, fragments or mutants thereof) or other ASLV RTs or subunits thereof (or derivatives, variants, fragments or mutants thereof). Particularly preferred AMV RTs and RSV RTs and other ASLV RTs include those provided by the present invention and described in detail above. More particularly preferred are those AMV RTs and RSV RTs having the genotype AMV α H/ β H⁺ RT or RSV α H/ β H⁺ RT. Such constructs are preferably made by mutating or modifying the gene encoding the α subunit to reduce or substantially reduce RNase H activity while the gene encoding the β subunit is not so mutated

or modified. The resulting polypeptides (produced by co-expression) will be reduced or substantially reduced in RNase H activity.

Other methods of cDNA synthesis which may advantageously use the present invention will be readily apparent to one of ordinary skill in the art.

5 Having obtained cDNA molecules or libraries according to the present methods, these cDNAs may be isolated for further analysis or manipulation. Detailed methodologies for purification of cDNAs are taught in the GENETRAPPER™ manual (Life Technologies, Inc.; Rockville, Maryland), which is incorporated herein by reference in its entirety, although alternative standard
10 techniques of cDNA isolation such as those described in the Examples below or others that are known in the art (*see, e.g., Sambrook, J., et al., Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, pp. 8.60-8.63 (1989)) may also be used.

15 In other aspects of the invention, the invention may be used in methods for amplifying and sequencing nucleic acid molecules. Nucleic acid amplification methods according to this aspect of the invention may be one- step (*e.g., one-step RT-PCR*) or two-step (*e.g., two-step RT-PCR*) reactions. According to the invention, one-step RT-PCR type reactions may be accomplished in one tube thereby lowering the possibility of contamination. Such one-step reactions
20 comprise (a) mixing a nucleic acid template (*e.g., mRNA*) with two or more polypeptides having reverse transcriptase activity and with one or more DNA polymerases and (b) incubating the mixture under conditions sufficient to amplify a nucleic acid molecule complementary to all or a portion of the template. Alternatively, amplification may be accomplished by mixing a template with two
25 or more polypeptides having reverse transcriptase activity (and optionally having DNA polymerase activity). Incubating such a reaction mixture under appropriate conditions allows amplification of a nucleic acid molecule complementary to all or a portion of the template. Such amplification may be accomplished by the reverse transcriptase activity alone or in combination with the DNA polymerase

activity. Two-step RT-PCR reactions may be accomplished in two separate steps. Such a method comprises (a) mixing a nucleic acid template (*e.g.*, mRNA) with two or more reverse transcriptases, (b) incubating the mixture under conditions sufficient to make a nucleic acid molecule (*e.g.*, a DNA molecule) complementary to all or a portion of the template, (c) mixing the nucleic acid molecule with one or more DNA polymerases and (d) incubating the mixture of step (c) under conditions sufficient to amplify the nucleic acid molecule. For amplification of long nucleic acid molecules (*i.e.*, greater than about 3-5 Kb in length), a combination of DNA polymerases may be used, such as one DNA polymerase having 3' exonuclease activity and another DNA polymerase being substantially reduced in 3' exonuclease activity. An alternative two-step procedure comprises the use of two or more polypeptides having reverse transcriptase activity and DNA polymerase activity (*e.g.*, *Tth*, *Tma* or *Tne* DNA polymerases and the like) rather than separate addition of a reverse transcriptase and a DNA polymerase.

Nucleic acid sequencing methods according to this aspect of the invention may comprise both cycle sequencing (sequencing in combination with amplification) and standard sequencing reactions. The sequencing method of the invention thus comprises (a) mixing a nucleic acid molecule to be sequenced with one or more primers, two or more reverse transcriptases, one or more nucleotides and one or more terminating agents, (b) incubating the mixture under conditions sufficient to synthesize a population of nucleic acid molecules complementary to all or a portion of the molecule to be sequenced, and (c) separating the population to determine the nucleotide sequence of all or a portion of the molecule to be sequenced. According to the invention, one or more DNA polymerases (preferably thermostable DNA polymerases) may be used in combination with or separate from the reverse transcriptases.

Amplification methods which may be used in accordance with the present invention include PCR (U.S. Patent Nos. 4,683,195 and 4,683,202), Strand Displacement Amplification (SDA; U.S. Patent No. 5,455,166; EP 0 684 315),

and Nucleic Acid Sequence-Based Amplification (NASBA; U.S. Patent No. 5,409,818; EP 0 329 822). Nucleic acid sequencing techniques which may employ the present compositions include dideoxy sequencing methods such as those disclosed in U.S. Patent Nos. 4,962,022 and 5,498,523, as well as more complex PCR-based nucleic acid fingerprinting techniques such as Random Amplified Polymorphic DNA (RAPD) analysis (Williams, J.G.K., *et al.*, *Nucl. Acids Res.* 18(22):6531-6535, 1990), Arbitrarily Primed PCR (AP-PCR; Welsh, J., and McClelland, M., *Nucl. Acids Res.* 18(24):7213-7218, 1990), DNA Amplification Fingerprinting (DAF; Caetano-Anollés *et al.*, *Bio/Technology* 9:553-557, 1991), microsatellite PCR or Directed Amplification of Minisatellite-region DNA (DAMD; Heath, D.D., *et al.*, *Nucl. Acids Res.* 21(24): 5782-5785, 1993), and Amplification Fragment Length Polymorphism (AFLP) analysis (EP 0 534 858; Vos, P., *et al.*, *Nucl. Acids Res.* 23(21):4407-4414, 1995; Lin, J.J., and Kuo, J., *FOCUS* 17(2):66-70, 1995). In a particularly preferred aspects, the invention may be used in methods of amplifying or sequencing a nucleic acid molecule comprising one or more polymerase chain reactions (PCRs), such as any of the PCR-based methods described above.

Kits

In another embodiment, the present invention may be assembled into kits for use in reverse transcription or amplification of a nucleic acid molecule, or into kits for use in sequencing of a nucleic acid molecule. Kits according to this aspect of the invention comprise a carrier means, such as a box, carton, tube or the like, having in close confinement therein one or more container means, such as vials, tubes, ampules, bottles and the like, wherein a first container means contains one or more polypeptides having reverse transcriptase activity. These polypeptides having reverse transcriptase activity may be in a single container as mixtures of two or more polypeptides, or in separate containers. The kits of the invention may

also comprise (in the same or separate containers) one or more DNA polymerases, a suitable buffer, one or more nucleotides and/or one or more primers.

5 In a specific aspect of the invention, the reverse transcription and amplification kits may comprise one or more components (in mixtures or separately) including one or more, preferably two or more, polypeptides having reverse transcriptase activity of the invention, one or more nucleotides needed for synthesis of a nucleic acid molecule, and/or a primer (*e.g.*, oligo(dT) for reverse transcription). Such reverse transcription and amplification kits may further comprise one or more DNA polymerases. Sequencing kits of the invention may
10 comprise one or more, preferably two or more, polypeptides having reverse transcriptase activity of the invention, and optionally one or more DNA polymerases, one or more terminating agents (*e.g.*, dideoxynucleoside triphosphate molecules) needed for sequencing of a nucleic acid molecule, one or more nucleotides and/or one or more primers. Preferred polypeptides having reverse transcriptase activity, DNA polymerases, nucleotides, primers and other components suitable for use in the reverse transcription, amplification and sequencing kits of the invention include those described above. The kits encompassed by this aspect of the present invention may further comprise additional reagents and compounds necessary for carrying out standard nucleic
15 acid reverse transcription, amplification or sequencing protocols. Such polypeptides having reverse transcriptase activity of the invention, DNA polymerases, nucleotides, primers, and additional reagents, components or compounds may be contained in one or more containers, and may be contained in such containers in a mixture of two or more of the above-noted components or
20 may be contained in the kits of the invention in separate containers.
25

Use of Nucleic Acid Molecules

5 The nucleic acid molecules or cDNA libraries prepared by the methods of the present invention may be further characterized, for example by cloning and sequencing (*i.e.*, determining the nucleotide sequence of the nucleic acid molecule), by the sequencing methods of the invention or by others that are standard in the art (*see, e.g.*, U.S. Patent Nos. 4,962,022 and 5,498,523, which are directed to methods of DNA sequencing). Alternatively, these nucleic acid molecules may be used for the manufacture of various materials in industrial processes, such as hybridization probes by methods that are well-known in the art. 10 Production of hybridization probes from cDNAs will, for example, provide the ability for those in the medical field to examine a patient's cells or tissues for the presence of a particular genetic marker such as a marker of cancer, of an infectious or genetic disease, or a marker of embryonic development. Furthermore, such hybridization probes can be used to isolate DNA fragments from genomic DNA or cDNA libraries prepared from a different cell, tissue or organism for further characterization. 15

20 The nucleic acid molecules of the present invention may also be used to prepare compositions for use in recombinant DNA methodologies. Accordingly, the present invention relates to recombinant vectors which comprise the cDNA or amplified nucleic acid molecules of the present invention, to host cells which are genetically engineered with the recombinant vectors, to methods for the production of a recombinant polypeptide using these vectors and host cells, and to recombinant polypeptides produced using these methods.

25 Recombinant vectors may be produced according to this aspect of the invention by inserting, using methods that are well-known in the art, one or more of the cDNA molecules or amplified nucleic acid molecules prepared according to the present methods into a vector. The vector used in this aspect of the invention may be, for example, a phage or a plasmid, and is preferably a plasmid. Preferred are vectors comprising *cis*-acting control regions to the nucleic acid

encoding the polypeptide of interest. Appropriate *trans*-acting factors may be supplied by the host, supplied by a complementing vector or supplied by the vector itself upon introduction into the host.

5 In certain preferred embodiments in this regard, the vectors provide for specific expression (and are therefore termed "expression vectors"), which may be inducible and/or cell type-specific. Particularly preferred among such vectors are those inducible by environmental factors that are easy to manipulate, such as temperature and nutrient additives.

10 Expression vectors useful in the present invention include chromosomal-, episomal- and virus-derived vectors, e.g., vectors derived from bacterial plasmids or bacteriophages, and vectors derived from combinations thereof, such as cosmids and phagemids, and will preferably include at least one selectable marker such as a tetracycline or ampicillin resistance gene for culturing in a bacterial host cell. Prior to insertion into such an expression vector, the cDNA or amplified nucleic acid molecules of the invention should be operatively linked to an appropriate promoter, such as the phage lambda P_L promoter, the *E. coli lac*, *trp* and *tac* promoters. Other suitable promoters will be known to the skilled artisan.

15 Among vectors preferred for use in the present invention include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; pcDNA3 available from Invitrogen; pGEX, pTrxfus, pTrc99a, pET-5, pET-9, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia; and pSPORT1, pSPORT2 and pSV•SPORT1, available from Life Technologies, Inc. Other suitable vectors will be readily apparent to the skilled artisan.

20
25 The invention also provides methods of producing a recombinant host cell comprising the cDNA molecules, amplified nucleic acid molecules or recombinant vectors of the invention, as well as host cells produced by such methods. Representative host cells (prokaryotic or eukaryotic) that may be produced according to the invention include, but are not limited to, bacterial cells, yeast

cells, plant cells and animal cells. Preferred bacterial host cells include *Escherichia coli* cells (most particularly *E. coli* strains DH10B and Stbl2, which are available commercially (Life Technologies, Inc; Rockville, Maryland)), *Bacillus subtilis* cells, *Bacillus megaterium* cells, *Streptomyces* spp. cells, *Erwinia* spp. cells, *Klebsiella* spp. cells and *Salmonella typhimurium* cells. Preferred animal host cells include insect cells (most particularly *Spodoptera frugiperda* Sf9 and Sf21 cells and *Trichoplusa* High-Five cells) and mammalian cells (most particularly CHO, COS, VERO, BHK and human cells). Such host cells may be prepared by well-known transformation, electroporation or transfection techniques that will be familiar to one of ordinary skill in the art.

In addition, the invention provides methods for producing a recombinant polypeptide, and polypeptides produced by these methods. According to this aspect of the invention, a recombinant polypeptide may be produced by culturing any of the above recombinant host cells under conditions favoring production of a polypeptide therefrom, and isolation of the polypeptide. Methods for culturing recombinant host cells, and for production and isolation of polypeptides therefrom, are well-known to one of ordinary skill in the art.

It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

Example 1: Cloning and Expression of RSV RNase H⁻ RT

General Methods

RSV H⁻ RT is a cloned form of retrovirus RT, in which both the α and the β subunits are mutated by a single amino acid change to eliminate RNase H activity. An RSV RT exhibiting substantially reduced RNase H activity is also produced when only the α subunit is mutated to substantially reduce RNase H activity (with the β subunit not being mutated in the RNase H domain). Mutations and plasmid constructions were conducted using standard molecular biology methods (*see, e.g.,* Sambrook, J., *et al., Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, NY: Laboratory Press (1989)), modified as described below.

Plasmid preparation. Plasmid preparations from 1 ml *E. coli* cultures were made by the alkaline lysis procedure (Sambrook, J., *et al., Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Laboratory Press (1989)). From 10 ml cultures, the preparation was additionally treated with phenol-chloroform, precipitated with ethanol, and resuspended in 50 μ l of Tris-EDTA (TE) buffer (20 mM Tris-HCl, 1 mM EDTA, pH 8.0). For bacmid preparations, care was taken to avoid shear of the DNA during handling (*i.e.,* no vortexing; phenol-chloroform extractions were gently rolled rather than shaken; slow pipetting).

PCR. Polymerase chain reactions were carried out in a Perkin-Elmer 9600 thermocycler. Reaction mixtures (50 μ l each) contained 0.5 units of *Taq* DNA polymerase, 1 μ M of each oligonucleotide, 50 μ M each of dCTP, dGTP, dTTP, and dATP, and about 100 ng of target DNA in a reaction buffer consisting of 50 mM KCl, 20 mM Tris-HCl (pH 8.3) and 5 mM MgCl₂. Unless otherwise noted, the cycling conditions for each PCR were 5 minutes at 94°C, followed by eight cycles of 15 seconds at 55°C/30 seconds at 72°C/15 seconds at 94°C, and then 1 minute at 72°C.

Gel electrophoresis and DNA fragment isolation and cloning. DNA was electrophoresed at about 10 V/cm in agarose gels in Tris-acetate-EDTA buffer (Sambrook, J., *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Laboratory Press (1989)) containing about 0.3 µg/ml ethidium bromide. Fragments were visualized with ultraviolet light and isolated from gel slices by the GlassMAX method (Simms, D., *et al.*, *Focus* 13:99 (1991)). DNA ligation reactions were performed using T4 DNA ligase under standard conditions (King, P.W., and Blakesley, R.W., *Focus* 8:1 (1986)), and *E. coli* DH10B cells were transformed by a modification of the CaCl₂ method (Lorow, D., and Jessee, J., *Focus* 12:19 (1990)).

Insect cell culture and baculovirus production. Samples (1 ml) of Sf21 insect cells at 5 x 10⁵ cells/ml were transfected with a mixture of 1 µg of bacmid DNA and 8 µl of Cellfectin in 0.2 ml SF900-II serum-free insect cell culture medium (Life Technologies, Inc.; Rockville, Maryland; *see* Godwin, G., and Whitford, W., *Focus* 15:44 (1993)), according to published procedures (Anderson, D., *et al.*, *Focus* 16:53 (1995)). For growth and propagation, Sf9 and Sf21 insect cells were passaged in SF900-II medium at 27°C in a shaking incubator at 100 rpm (for 600 ml cultures) or 130 rpm (for all other cultures). Care was taken to avoid allowing the cultures to exceed 4 x 10⁶ cells/ml during growth or to drop below 0.5 x 10⁶ cells/ml during dilution. To expand viral populations, Sf21 cells at about 1 X 10⁶ cells/ml were infected with enough viral stock (about 0.2% (v/v) virus/culture) to allow growth to about 2 x 10⁶ cells/ml, but not more than 4 x 10⁶ cells/ml. After 72 hours, the culture was centrifuged (2,000 rpm for 10 min) and the supernatant was decanted and stored in the dark at 4°C. For infection of cells for protein production, Sf21 insect cells at about 1.5 x 10⁶ cells/ml were infected with enough virus to allow no growth or growth to less than 2.5 x 10⁶ cells/ml. After 72 hours, the culture was harvested by

centrifugation at 1,000 rpm for 5 minutes and cells were resuspended in 15 ml of PBS (0.2 g/liter KCl, 0.2 g/liter KH₂PO₄, 8 g/liter NaCl, 1.15 g/liter Na₂HPO₄, 2.16 g/liter Na₂HPO₄•7H₂O) per 500 ml culture.

Cloning and Expression of Genes Encoding the RSV RT α and β Subunits

Both the α and β subunits of RSV RT are produced by proteolytic processing of larger polypeptide precursors (Gerard, G. F., in: *Enzymes of Nucleic Acid Synthesis and Modification, Vol. I: DNA Enzymes*, Jacob, S.T., ed., Boca Raton, FL: CRC Press, pp. 1-38 (1983)). To obviate the requirement for proteolytic processing, the coding sequence for RSV RT was mutagenized and subcloned such that both the α and β subunits were encoded by genes with standard start and stop translational signals. Both genes were mutagenized in the RNase H region, although construction of any combination of subunits (e.g., α RNase H⁻/ β RNase H⁺; α RNase H⁺/ β RNase H⁺; α RNase H⁺/ β RNase H⁻; α RNase H⁻/ β RNase H⁻) may be accomplished in this same manner. It has been discovered that RSV RT α RNase H⁻/ β RNase H⁺ is substantially reduced in RNase H activity (approximately 5% of wildtype). A sequence encoding an affinity tag was added to the carboxy end of the β subunit.

Mutagenesis and subcloning of the amino end, the carboxy end and the middle of the RSV RT β subunit. The RSV RT gene was mutagenized to add an ATG codon and an NdeI site to the amino end of the sequence coding for the mature RT polypeptide. This mutagenesis was accomplished by PCR using a pJD100 target (Figures 1, 7) (Wilkerson, V.W., et al., *J. Virol.* 55:314-321 (1985)) and the following oligonucleotides:

AUG GAG AUC UCU CAT ATG ACT GTT GCG CTA CAT CTG GCT
(SEQ ID NO:1)

AAC GCG UAC UAG U GTT AAC AGC GCG CAA ATC ATG CAG
(SEQ ID NO:2).

PCR was performed, and PCR products purified, as described above and the PCR products were cloned into pAMP18 by UDG cloning (Buchman, G.W., *et al.*, *Focus* 15:36 (1993)) to form plasmid pAMP18N (Figures 1, 8).

Following mutagenesis and cloning of the amino end, the carboxy end of the gene for the β subunit of RSV was mutagenized and subcloned from PJD100 by PCR and UDG cloning into pAMP1 (Figure 1), using the following oligonucleotides:

CUA CUA CUA CUA GGT ACC CTC TCG AAA AGT TAA ACC
(SEQ ID NO:3)

CAU CAU CAU CAU CTC GAG TTA TGC AAA AAG AGG GCT CGC CTC
ATC (SEQ ID NO:4).

These oligonucleotides were designed to introduce a translational stop codon in the β gene at the site in which the "p4" region was cleaved from the β p4 polypeptide, and to add an *Xho*I site after the end of the gene. The PCR products were purified by gel electrophoresis and cloned into pAMP1 by UDG cloning, forming pAMP1C (Figures 1, 9). Note that this carboxy end is without a His tag, which was added later to form the final construct.

To add the middle region of the RSV RT β subunit, the 2.3 kb *Hpa*I-*Kpn*I fragment from pJD100 (Figure 1) that encodes the middle of the β subunit of RSV RT was cloned into the *Hpa*I-*Kpn*I sites of pAMP18N, forming pAMP18NM (Figures 1, 10). To add the carboxy end of the RSV RT β subunit, the 113 bp *Kpn*I-*Eco*RI fragment encoding the carboxy end of the β subunit gene was cloned from pAMP1C into the *Kpn*I-*Eco*RI sites of pAMP18NM, forming pAMP18B (Figures 2, 11).

Following construction of the RSV RT β subunit, which contains RNase H activity, this gene was mutagenized by site-directed mutagenesis to produce a construct encoding a RSV RT β subunit that is substantially reduced in RNase H activity (*i.e.*, "RNase H"). A 3 Kb *Pst*I fragment from pJD100 (Figure 2)

containing the entire RT gene was cloned into M13mp19, forming M13RT (Figures 2, 12). Single-stranded DNA containing uracil was isolated from *E. coli* strain CJ236 (Bio-Rad; Hercules, CA; and Cathy Joyce, Yale University, New Haven, CT) after infection with M13RT phage containing the *Pst*I fragment. To mutate the RNase H region and to introduce an *Sst*II site, the following oligonucleotide was used:

GGA CCC ACT GTC TTT ACC GCG GCC TCC TCA AGC ACC
(SEQ ID NO:5)

This oligonucleotide induced the substitution of an alanine residue in place of the aspartate residue at position 450 of the RT, forming M13RTH⁻ (Figures 2, 13). In an alternative approach to generate a RNase H⁻ RSV RT, Glu484 may be mutated to glutamine and/or Asp505 may be mutated to asparagine. To convert the β subunit back to RNase H⁺, an oligonucleotide primer having the wildtype sequence may be used. Alternatively, deletions or insertions can be made in the RNase H region to substantially reduce RNase H activity. DNA sequencing was used to confirm the Asp450→Ala450 mutation, and the 426 bp *Bgl*III-*Bst*EII fragment from M13RTH⁻ was cloned into the *Bgl*III-*Bst*EII sites of pAMP18B, replacing the RNase H region and forming pAMP18BH⁻ (Figures 2, 14).

Mutagenesis and subcloning of the gene encoding the RSV RT α subunit.

To create a gene which codes for the α subunit of RSV RT, two oligonucleotides were used to mutagenize the amino end of the RNase H⁻ mutant RSV RT gene from pDBH⁻ (Figures 3, 15) and to introduce a translational stop codon where avian retroviral protease p15 normally cleaves the precursor polypeptide to make the α subunit:

CAU CAU CAU CAU CCC GGG TTA ATA CGC TTG GAA GGT GGC
(SEQ ID NO:6)

CUA CUA CUA CUA TCA TGA CTG TTG CGC TAC ATC TG
(SEQ ID NO:7)

PCR cycling conditions were 5 minutes at 94°C, followed by 8 cycles of
15 seconds at 55°C/2 minutes at 72°C/15 seconds at 94°C, and then 2 minutes
at 72°C. The PCR products were cloned into pAMP1 by UDG cloning as
described above, forming pAMP1A (Figures 3, 16).

Addition of a His₆ tag to the carboxy end of the RSV β subunit. A His₆ tag
was added to the carboxy end of the RSV RT β subunit by site-directed
mutagenesis, and the mutant sequence was subcloned from pJD100 by PCR and
UDG cloning as described above, using the following oligonucleotides:

CUA CUA CUA CUA GGT ACC CTC TCG AAA AGT TAA
(SEQ ID NO:8)

CAU CAU CAU CAU GAG GAA TTC AGT GAT GGT GAT GGT GAT GTG
CAA AAAG AGG
(SEQ ID NO:9)

These oligonucleotides were designed to introduce a translational stop codon in
the gene, and to add a histidine tag to the end of the protein. The gene product
was thus a polypeptide to which 6 histidines were added to the carboxy end. The
PCR products were purified by gel electrophoresis and inserted into pAMP1 by
UDG cloning, forming pAMPChis.

To remove the carboxy end of the RSV RT β gene and replace it with the
carboxy end containing a His₆ tag, the number of *Kpn*I sites in the baculovirus
vector containing the RSV RT β gene had to be reduced. Plasmid pDBH-
(Figure 4) was cleaved with *Afl*III, and the site was "blunted" with the Klenow
fragment of *E. coli* DNA polymerase I, the polymerase was inactivated by heat
treatment, and the vector was further cleaved with *Pvu*II (Figure 4). The deleted
vector (5.9 kb) was then purified by gel electrophoresis, ligated to itself and used
to transform *E. coli* strain DH10B, forming pDBH-Kpn (Figures 4, 17). The

KpnI-SstI fragment with the carboxy end of β with the His₆ tag was cloned from pAMPChis into the *KpnI-SstI* sites of pDBH-Kpn, forming pDBH-KpnHis (Figures 4, 18).

Cloning of RSV α and β subunits into a baculovirus expression vector.

During the course of this work, a construct was made in which the RSV RT β gene was to be expressed in a baculovirus vector with a His₆ tag at the amino end. However, a mutation introduced during the construction caused the reading frame of the tag to be different from that of the β gene. Since portions of this construct were used to construct the final baculovirus expression vector, its construction is described here.

To introduce a His₆ tag and an *NdeI* site into the baculovirus vector, the following oligonucleotides were annealed and cloned into pFastBac Dual (Harris, R., and Polayes, D.A., *Focus* 19:6-8 (1997); Figure 5):

ACTG GAA TTC ATG CCA ATC CAT CAC CAT CAC CAT CAC CCG T
(SEQ ID NO:10)

ACGT GTC GAC CAT ATG GAT GAC TAG GTG AAA CGG GTG ATG G
(SEQ ID NO:11)

Both oligonucleotides were formulated into TE buffer at a concentration of 100 μ M, and 5 μ l of each oligonucleotide formulation were mixed with 15 μ l of water and 2 μ l of 10X React2 buffer (500 mM NaCl, 500 mM Tris-HCl, 100 mM MgCl₂, pH 8.0) in a single tube. This tube was heated in a 65°C water bath and cooled slowly over 60 minutes to 25°C. The resulting product was a double-stranded DNA molecule with an *EcoRI* site at the 5' end (underlined), and a *SalI* site at the 3' end (*italicized*):

5'-ACTG GAA TTC ATG CCA ATC CAT CAC CAT CAC CAT CAC CCG
TTT CAC CTA GTC ATC CAT ATG *GTC GAC* ACGT-3' (SEQ ID NO:12)

This product was cleaved with *EcoRI* and *SalI* and the desired fragment was cloned into the *EcoRI-SalI* sites of vector pFastBac Dual by standard techniques (Harris, R., and Polayes, D. A., *Focus* 19:6-8 (1997)), resulting in the formation of plasmid pFastBac Dual Nde (Figure 5). The *NdeI-XhoI* B fragment of pAMP18BH⁺ was cloned into the *NdeI-SalI* sites of pFastBac Dual Nde to create pDBH⁺ (Figures 5, 15).

To clone the RSV RT α gene into a baculovirus vector, the α gene was excised from pAMP1A with *SmaI* and *BspHI* and subcloned into the *NcoI-SmaI* sites of pFastBac Dual, creating pDA (Figures 5, 19). This placed the RSV RT α gene downstream from the baculovirus P10 promoter. The RSV α peptide gene and P10 promoter were excised with *RsrII* and *SmaI*, and cloned into the *RsrII-SmaI* sites of pDBH⁺, forming pDABH⁺ (Figures 4, 20).

To replace the RSV RT β gene in pDABH⁺ with the carboxy His₆-tagged β gene in pDBH-KpnHis, pDBH-KpnHis was cleaved with *NdeI*, the site blunt-ended with Klenow fragment, and the β gene with the carboxy His₆ tag was then released with *SstI* (Figure 6). pDABH⁺ was cleaved with *EcoRI*, the site blunt-ended with Klenow fragment, and then the β gene was removed by digesting with *SstI*. The vector fragment, which was a blunt-end *SstI* fragment of pFastBac Dual with the α gene cloned in front of the p10 promoter, was ligated to the blunt-end *SstI* fragment with the carboxy His-tagged β gene from pDBH-KpnHis. *E. coli* DH10B cells were transformed with this construct, and a transformant containing pDABH-His (Figures 6, 21) was selected.

The recombinant host cell comprising plasmid pDABH-His, *E. coli* DH10B(pDABH-His), was deposited on April 15, 1997, with the Collection, Agricultural Research Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois 61604 USA, as Deposit No. NRRL B-21679.

Using the deposited plasmid, one of ordinary skill in the art may easily produce, using standard genetic engineering techniques (such as those for site-directed mutagenesis described above), plasmids encoding various forms of the α

and/or β subunits of RSV RT (e.g., α RNase H⁺/ β RNase H⁺; α RNase H⁻/ β RNase H⁻; α RNase H⁺/ β RNase H⁻; and α RNase H⁻/ β RNase H⁺).

Transfection of insect cells for virus and RSV RT production. To prepare vectors for the transfection of insect cells, it was first necessary to insert the RSV RT gene constructs into a baculovirus genome. One method of accomplishing this insertion is by using the site-specific transposon Tn7 (Luckow, V.A., in: *Recombinant DNA Technology and Applications*, Prokop, A., *et al.*, eds., New York: McGraw-Hill (1991)). In DH10Bac host cells, most of the baculovirus genome is represented on a low copy plasmid (a "bacmid") which also contains a Tn7 insertion site within a gene (Harris, R., and Polayes, D. A., *Focus* 19:6-8 (1997)). Transposition of Tn7 is facilitated by the Tn7 transposase, which is produced from a second plasmid in DH10Bac host cells, and pFastBac DUAL is constructed with the "right" and "left" regions of Tn7 flanking the promoter and cloning sites.

To insert the present constructs into a bacmid expression vector, pDABH-His was transformed into DH10Bac cells and transposition of sequences encoding the RSV RT β gene (whose synthesis is directed by the baculovirus polyhedrin promoter) and the RSV RT α gene (whose synthesis is directed by the baculovirus P10 promoter) was followed by screening for loss of β -galactosidase activity following transposition to the site on the bacmid which codes for the β -galactosidase α peptide (Harris, R., and Polayes, D. A., *Focus* 19:6-8 (1997)). Bacmid DNA was then prepared from 10 ml transformant cultures by a slight modification of a standard miniprep procedure (Sambrook, J., *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989)). About 1 μ g of the bacmid DNA was then used to transfect Sf21 insect cells using the cationic lipid Cellfectin (Anderson, D., *et al.*, *Focus* 16:53 (1995)).

For expansion of primary virus, the supernatant was removed from the transfected cells about 72 hours after transfection and 1 ml was used to infect

35 ml of *Sf*21 insect cells (about 1.2×10^5 cells/ml). After 72 hours, the culture was centrifuged (2,000 rpm; 10 min) and the supernatant was decanted and used as a secondary viral stock. The secondary virus stock was expanded similarly by infecting 35 ml of *Sf*21 cells with 0.1 ml of the secondary stock.

5 Virus stocks were then used to infect *Sf*21 cells for the expression of RSV RT. In preliminary experiments, expression of RT activity from infected cells was found to be maximal about 72 hours after infection. For test expressions, 70 ml of *Sf*21 cells were infected with 5 ml of the virus stock, and the cells were harvested 72 hours after infection by centrifugation at 1,000 rpm for 5 minutes, resuspension in PBS (2.5% of the culture volume) and recentrifugation at 1,000 rpm for 5 minutes. Supernatants were removed and the cells were stored at -70°C until use. For larger scale production, 600 ml of cells in 2.8 liter Fernbach flasks were infected with 2 ml of virus stock, and the cells were harvested 72 hours after infection.

15 **Example 2: Isolation of RSV RNase H⁻ RT**

To provide purified recombinant RSV H⁻ RT, cloned RSV H⁻ RT was overexpressed in cultured insect cells as described in Example 1 and purified by affinity and ion-exchange chromatography. The RSV RT produced comprises the α and β subunits. Isolation of RSV RT provides a substantially pure RSV RT in which contaminating enzymes and other proteins have been substantially removed, although such contaminants need not be completely removed.

20 **Buffers.** The pH of all buffers was determined at 23°C , and buffers were stored at 4°C until use. Crack Buffer contained 50 mM Tris-HCl (pH 7.9), 0.5 M KCl, 0.02% (v/v) Triton X-100 and 20% (v/v) glycerol. Just before use, the following protease inhibitors (Boehringer Mannheim; Indianapolis, IN) were added to Crack Buffer at the final concentrations indicated: leupeptin (2 $\mu\text{g/ml}$), Pefabloc (48 $\mu\text{g/ml}$), pepstatin A (2 $\mu\text{g/ml}$), benzamidine (800 $\mu\text{g/ml}$) and PMSF

(50 µg/ml). Buffer A contained 20 mM Tris-HCl (pH 7.9), 0.25 M KCl, 0.02% (v/v) Triton X-100, and 10% (v/v) glycerol. Buffer B was Buffer A with 1 M imidazole added. Buffer S contained 50 mM Tris-HCl (pH 8.2), 0.02% (v/v) Triton X-100, 10% (v/v) glycerol, 0.1 mM EDTA and 1 mM dithiothreitol (DTT). Buffer T was Buffer S with 1 M KCl added. Buffer H contained 20 mM potassium phosphate (pH 7.1), 0.02% (v/v) Triton X-100, 20% (v/v) glycerol, 0.1 mM EDTA and 1 mM DTT. Buffer J was Buffer H with 1 M KCl added. Storage Buffer contained 200 mM potassium phosphate (pH 7.1), 0.05 % (v/v) NP-40, 50% glycerol (v/v), 0.1 mM EDTA, 1 mM DTT, and 10% (w/v) trehalose.

Extract preparation. Frozen insect cells (25 g) were thawed and a slurry prepared at 4°C with 50 ml of Crack Buffer plus inhibitors. Cells were disrupted at 4°C by sonication with a Fisher 550 Sonicator at 25% of maximum power. The disrupted crude extract was clarified by centrifugation at 27,000 x g for 30 minutes at 4°C.

RSVRT isolation. Following clarification, the extract was fractionated and RSV RT purified by column chromatography at 4°C. The clarified crude extract was loaded unto a 30 ml Chelating Sepharose Fast Flow column (Pharmacia; Piscataway, NJ) charged with NiSO₄ as per manufacturer's instructions and equilibrated in 99.5% Buffer A + 0.5% Buffer B. The column was washed with one column volume of 99% Buffer A + 1% Buffer B and then with 10 column volumes of 98.5% Buffer A + 1.5% Buffer B. RT was eluted with a 10-column volume linear gradient of 98.5% Buffer A + 1.5% Buffer B to 75% Buffer A + 25% Buffer B.

During purification, reverse transcriptase activity was assayed with poly(C)-oligo(dG), which is specific for reverse transcriptase (Gerard G. F., *et al.*, *Biochemistry* 13:1632-1641 (1974)). RT unit activity was defined and assayed as described (Houts, G. E., *et al.*, *J. Virol.* 29:517-522 (1979)). Using the poly(C)-oligo(dG) assay, the peak fractions of RT activity from the Chelating Sepharose Fast Flow column (10 to 17% Buffer B) were pooled, diluted with an equal

volume of Buffer S, and loaded on a 5 ml AF-Heparin-650M column (TosoHaas; Montgomeryville, PA) equilibrated in Buffer S. After a wash with 12 column volumes of 90% Buffer S + 10% Buffer T, the column was eluted with a 15-column volume linear gradient of Buffer S to 30% Buffer S + 70% Buffer T. The peak fractions of RT activity (43 to 50% Buffer T) were pooled, diluted with 2.5 volumes of Buffer H, and loaded onto a Mono S HR 5/5 column (Pharmacia; Piscataway, NJ) equilibrated in Buffer H. After a wash with 20 column volumes of 85% Buffer H + 15% Buffer J, the column was eluted with a 20 column volume gradient of 85% Buffer H + 15% Buffer J to 50% Buffer H + 50% Buffer J. The RT peak fractions were pooled, dialyzed against Storage Buffer overnight, and stored at -20°C.

Following purification, RSV H⁻ RT was found to be >95% homogeneous as judged by SDS-PAGE. The purified enzyme was also found to be substantially lacking in RNase and DNase contamination, and substantially reduced in RNase H activity.

Example 3: Preparation of Full-length cDNA Molecules

Enzymes. SuperScript II RT (SS II RT), a cloned form of Moloney murine leukemia virus (M-MLV) RT lacking demonstrable RNase H activity (*i.e.*, an "RNase H⁻ RT"), was from Life Technologies, Inc. (Rockville, Maryland). M-MLV RT, a cloned murine RT with full RNase H activity (*i.e.*, an "RNase H⁺ RT"), was also from Life Technologies, Inc. AMV RT, an RNase H⁺ uncloned form of avian myeloblastosis virus RT, was from Seikagaku America, Inc. RSV H⁻ RT was prepared as described in Examples 1 and 2. Recombinant *Tth* DNA polymerase, a cloned, thermophilic DNA polymerase from *Thermus thermophilus* with reverse transcriptase activity, was from Perkin Elmer.

Synthetic mRNA. A 7.5 kilobase (Kb) synthetic mRNA with a 120-nucleotide 3' poly(A) tail (Life Technologies, Inc.; Rockville, Maryland) was used

as template to test the efficiency of various enzymes alone or in combination.

cDNA Synthesis Reaction Mixtures. Reaction mixtures (20 μ l each) contained the following components unless specified otherwise: 50 mM Tris-HCl (pH 8.4 at 24°C), 75 mM KCl, 10 mM dithiothreitol, 1 mM each of [³²P]dCTP (300 cpm/pmole), dGTP, dTTP, and dATP, 25 μ g/ml (p(dT)₂₅₋₃₀), 125 μ g/ml of 7.5 Kb mRNA, and 35 units of cloned rat RNase inhibitor. Reaction mixtures with RTs alone or in combination contained the following:

SS II RT alone:	0.5 mM dNTPs, 3 mM MgCl ₂ , and 200 units of SS II RT
RSV H ⁻ RT alone:	7.5 mM MgCl ₂ and 7 units of RSV H ⁻ RT;
AMV RT alone:	50 mM KCl, 10 mM MgCl ₂ , 4 mM sodium pyrophosphate and 14 units of AMV RT
SS II RT plus RSV H ⁻ RT:	5 mM MgCl ₂ , 200 units of SS II RT, and 7 units of RSV H ⁻ RT;
<i>Tth</i> DNA Polymerase alone:	0.5 mM dNTPs, 1 mM MnCl ₂ and 5 units of <i>Tth</i> DNA Polymerase
M-MLV H ⁺ RT alone:	0.5 mM dNTPs, 3 mM MgCl ₂ , 50 μ g/ml actinomycin D and 200 units of M-MLV H ⁺ RT
<i>Tth</i> DNA Polymerase plus either SS II RT or RSV H ⁻ RT:	same as <i>Tth</i> DNA Polymerase alone, plus either 200 units of SS II RT or 7 units of RSV H ⁻ RT
M-MLV H ⁺ RT plus SS II RT:	same as M-MLV H ⁺ RT alone, plus 200 units of SS II RT;
M-MLV H ⁺ plus either AMV RT or RSV H ⁻ RT:	5 mM MgCl ₂ , 50 μ g/ml actinomycin D, 200 units of M-MLV H ⁺ RT, and either 14 units of AMV RT or 7 units of RSV H ⁻ RT
AMV RT plus either SS II RT or RSV H ⁻ RT:	5 mM MgCl ₂ , 50 μ g/ml actinomycin D, 14 units of AMV RT, and either 200 units of SS II RT or 7 units of RSV H ⁻ RT

When RTs were used in combination, one enzyme was added first followed immediately by the addition of an aliquot of the second enzyme. In some cases in which a single enzyme was used, a second aliquot of the same enzyme was added as a control to assess the effect of doubling the amount of the single enzyme.

5 All cDNA synthesis reactions were carried out at 45°C for 50 minutes, and the resultant cDNA product was detectably labeled by the RT-catalyzed incorporation of a ³²P-labeled deoxyribonucleoside triphosphate precursor. The total yield of cDNA was determined by acid precipitation of a portion of the cDNA product and counting it in a scintillation counter. The ³²P-labeled cDNA product in the remainder of the reaction mixture was fractionated by alkaline agarose gel electrophoresis (Carmichael, G.G., and McMaster, G.K., *Meth. Enzymol.* 65:380-385 (1980)). The gel was dried and the size distribution of the cDNA product was established by autoradiography. Using the autoradiographic film as a template, the dried gel was cut and analyzed by scintillation counting to establish the fraction of full length (7.5 Kb) product synthesized.

10
15 Tables 1 and 2 show the total amount of cDNA synthesized and full length cDNA synthesized, respectively, from the 7.5 Kb mRNA by enzymes alone or in combination. The following conclusions can be drawn:

20 1. When RTs were present alone, the highest yields of total and full length product were obtained with the RNase H⁻ forms of RT. With either RSV H⁻ RT or SS II RT, the total yield was almost double the highest yield obtained with an RNase H⁺ enzyme (1011 and 946 ng, respectively, versus 607 ng). The effect of removing RNase H from the reaction was even more dramatic when full length yields were examined. In this case, yields were at least tripled (234 and 25 208 ng for RSV H⁻ and SS II RT versus 79 and 26 for M-MLV H⁺ RT and AMV RT, respectively). These results demonstrate the dramatic positive effect of eliminating RNase H from RT.

2. When RTs were combined, several effects were observed. Mixing RTs from different sources, whether RNase H⁻ or RNase H⁺, increased total and

full length yields. This is consistent with the hypothesis that pausing at sites unique to one enzyme can be reduced by a second RT with a different set of pause sites. However, by far the greatest yields of total and full length cDNA product were obtained when two different RNase H⁻ RTs were combined (see shaded boxes in Tables 1 and 2). These results indicate that the two RNase H⁻ enzymes cooperate to synthesize full-length cDNA molecules: the first enzyme synthesizes truncated cDNA molecules, which are then extended to full-length via the activity of the second enzyme. Thus, the compositions and methods of the present invention facilitate the synthesis of full-length cDNA molecules.

Table 1. Total Yield of cDNA Synthesized by Various Enzymes from 7.5-Kb mRNA ¹						
Enzyme Added Second	Enzyme Added First					
		RSV H ⁻ RT	SS II RT	M-MLV H ⁺ RT	AMV RT	<i>Tth</i> DNA Polymerase
	(None)	1,011	946	607	316	297
	RSV H ⁻ RT	952	1,286	802	479	-- ²
	SS II RT	1,362	1,046	751	827	--
	M-MLV H ⁺ RT	848	621	599	867	--
	AMV RT	533	922	802	353	--
	<i>Tth</i> DNA Polymerase	218	996	--	--	--

¹Mass (ng) of total cDNA product

²not tested

Table 2. Yield of Full Length cDNA Synthesized by Various Enzymes from 7.5 Kb mRNA¹

Enzyme Added Second	Enzyme Added First					
		RSV H ⁻ RT	SS II RT	M-MLV H ⁺ RT	AMV RT	<i>Tth</i> DNA Polymerase
	(None)	234	208	79	26	<1
	RSV H ⁻ RT	211	258	128	71	-- ²
	SS II RT	290	229	105	141	--
	M-MLV H ⁺ RT	128	79	69	96	--
	AMV RT	83	148	100	38	--
	<i>Tth</i> DNA Polymerase	5	176	--	--	--

¹Mass (ng) of full-length (7.5 kb) cDNA product

²not tested

In the mixing experiments summarized in Tables 1 and 2, two reverse transcriptases were added simultaneously to a reaction under conditions that may have been suboptimal for a single given RT. This was the case when an avian and a murine RT were used together, since the MgCl₂ concentration was set at 5 mM, between the optima of 3 mM and 7.5 mM for murine and avian RT, respectively. In addition, full advantage could not be taken in these experiments of the thermal stability of avian RNase H⁻ RT in reactions containing a less thermostable murine RT.

To use multiple enzymes and to address the fact that different enzymes may have different optimal conditions, sequential additions or separate use of the enzyme may be done in accordance with the methods of the invention. For example, cDNA could be synthesized from different aliquots of the same RNA in separate reaction tubes with different RTs under reaction conditions optimal for

each RT. Subsequently, the cDNAs from each reaction could be mixed before performing further manipulations. Alternatively, RTs could be used singly and sequentially in one tube to perform cDNA synthesis. That is, SuperScript II could be used first to copy an RNA population under optimal reaction conditions, and then conditions could be adjusted to optimal for RSV H⁻ RT in the same tube, and further synthesis could be performed with the avian RT at elevated temperature.

Example 4: Cloning and Expression of Avian Myeloblastosis Virus (AMV) RT and AMV RNase H⁻ (AMV H⁻) RT

General Methods

The AMV RT of the present invention is a cloned form of avian retrovirus RT. The AMV H⁻ RT is a variant of cloned AMV RT in which both the α and the β subunits are mutated by a single amino acid change to eliminate RNase H activity, although AMV RT substantially reduced in RNase H activity is also produced by mutating the α subunit alone (the β subunit not containing a mutation in the RNase H domain). Mutations and plasmid constructions were conducted using standard molecular biology methods (*see, e.g.,* Sambrook, J., *et al., Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, NY: Laboratory Press (1989)), modified as described below. Plasmid preparation, PCR, gel electrophoresis, DNA fragment isolation and cloning, insect cell culture and baculovirus production were all performed as described for RSV RT cloning and expression in Example 1.

Cloning and Expression of Genes Encoding the AMV RT α and β Subunits

To clone AMV RT, AMV viral RNA was prepared (Strauss, E.M., *et al., J. Virol. Meth. 1*:213 (1980)) from purified (Grandgenett, D.P., *et al., Appl. Microbiol. 26*:452 (1973)) AMV obtained from Life Sciences (St. Petersburg, Florida). AMV RT cDNA was prepared from AMV viral RNA with the

SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Life Technologies, Inc.; Rockville, Maryland) following the instructions in the kit manual. A primer specific for AMV RT was used that adds a *NotI* site to the cDNA. Following preparation, AMV RT cDNA was cloned into pSPORT1 that had been treated with *SaII* and *NotI*, resulting in a vector (pSPORT8) comprising the AMV RT gene (Figure 22).

Both the α and β subunits of AMV RT are produced by proteolytic processing of larger polypeptide precursors (Gerard, G.F., in: *Enzymes of Nucleic Acid Synthesis and Modification, Vol. I: DNA Enzymes*, Jacob, S.T., ed., Boca Raton, Florida: CRC Press, pp. 1-38 (1983)). To obviate the requirement for proteolytic processing, the coding sequence for AMV RT was mutagenized and subcloned such that both the α and β subunits were encoded by genes with standard start and stop translational signals. To make RNase H⁻ constructs, both α and β genes were mutagenized in the RNase H region, although construction of any combination of subunits (*e.g.*, α RNase H⁻/ β RNase H⁺; α RNase H⁺/ β RNase H⁺; α RNase H⁺/ β RNase H⁻; α RNase H⁻/ β RNase H⁻) may be accomplished in this same manner. It has been discovered that AMV RT α RNase H⁻/ β RNase H⁺ is substantially reduced in RNase H activity (approximately 5% of wildtype). A sequence encoding an affinity tag was added to the carboxy end of the β subunit.

Synthesis of cDNA from AMV RNA. The AMV RNA was copied into DNA using a 3' primer which is complementary to the 3' end of the AMV RT gene, and which adds a *NotI* site to the 3' end of the gene (Figure 22). The 5' end of the resulting cDNA was made into a *SaII* end by the addition of a *SaII* adapter. This cDNA was then cloned into a *SaII-NotI* cleaved vector (pSPORT1). First strand cDNA was synthesized using Superscript II RT (Gerard, G.F., *et al.*, *FOCUS 11*:66 (1989)) and a gene specific primer (Oligonucleotide #1) instead of the *NotI* primer-adapter:

Oligonucleotide #1 (SEQ ID NO:13):

5' GACTAGTTCTAGATCGCGAGCGGCCGCCCATTA ACTCTCGTTGG
CAGC 3'

The second strand synthesis was achieved using DNA polymerase I in combination with *E. coli* RNase H and DNA ligase at 16°C and subsequently polishing the termini with T4 DNA polymerase. The cDNA was deproteinized and precipitated with ethanol, and *Sal*I adapters consisting of Oligonucleotides #2 and #3 were ligated to the cDNA:

Oligonucleotide #2 (SEQ ID NO:14):

5' TCGACCCACGCGTCCG 3'

Oligonucleotide #3 (SEQ ID NO:15):

5' CGGACGCGTG GG 3'

The addition of the adapters was followed by digestion with *Not*I. Size fractionation of the cDNA was done on 1 ml prepacked columns provided with the cDNA cloning kit. The amount of cDNA in each fraction was calculated from the specific activity of incorporated ³²P label, and the size of the cDNA was determined by autoradiography of an agarose gel. Those fractions that were greater than 3 Kb were selected for cloning.

Cloning AMV cDNA into a Vector. The cDNA was ligated into *Sal*I-*Not*I cleaved pSPORT1 vector and then the ligated cDNA was used to transform *E. coli* MAX EFFICIENCY DH10B™ competent cells (Life Technologies, Inc., Rockville, Maryland). After transformation, aliquots of cells were plated on LB plates containing ampicillin. Twelve colonies were picked and 1 ml cultures were grown for mini-preps. Gels were run to check for certain fragments after digesting with restriction enzymes *Sal*I, *Mlu*I, *Pst*I, *Apa*I, *Dra*III, *Sph*I and *Bgl*II. One plasmid, pSPORT8, was selected since the insert was large enough to code for the

AMV RT gene (3 Kb) and a *Pst*I site was present which indicated that the 5' terminus of the AMV RT gene was present (Figure 22).

Mutagenesis and subcloning of the amino end, the carboxy end and the middle of the AMV RT β subunit. The AMV RT gene was mutagenized to add an ATG codon and an *Eco*RI site to the amino end of the sequence coding for the mature RT polypeptide by PCR with pSPORT8 as the target and the following oligonucleotides:

Oligonucleotide #4 (SEQ ID NO:16):

5' AUG GAG AUC UCU GAA TTC ATG ACT GTT GCG CTA CAT CTG
GCT 3'

Oligonucleotide #5 (SEQ ID NO:2):

5' AAC GCG UAC UAG U GTT AAC AGC GCG CAA ATC ATG CAG 3'

PCR was performed, and PCR products purified, as described above. The PCR reaction was treated with *Dpn*I to destroy the target and the PCR product was cloned into pAMP18 by UDG cloning (Buchman, G.W., *et al.*, *Focus* 15:36 (1993)), forming plasmid pAMVN (Figures 23, 26).

Following mutagenesis and cloning of the amino end, a His₆ affinity tag, a *Xho*I site and a translational stop codon were added to the carboxy end of the gene for the β subunit of AMV RT in pSPORT8 by PCR using the following oligonucleotides:

Oligonucleotide #6 (SEQ ID NO:3):

5' CUA CUA CUA CUA GGT ACC CTC TCG AAA AGT TAA ACC 3'

Oligonucleotide #7 (SEQ ID NO:9):

5' CAU CAU CAU CAU GAG GAA TTC AGT GAT GGT GAT GGT GAT
GTG CAA A AAG AGG 3'

PCR was performed, and PCR products purified, as described above. The PCR reaction was treated with *DpnI* to destroy the target and the PCR product was cloned into pAMP18 by UDG cloning (Buchman, G.W., *et al.*, *Focus* 15:36 (1993), forming plasmid pAMVC (Figures 23, 27).

To add the middle region of the AMV RT β subunit, the 2.3 Kb *HpaI*-*KpnI* fragment from pSPORT8 that encodes the middle of the β subunit of AMV RT was cloned into the *HpaI*-*KpnI* sites of pAMVN, forming pAMVNM (Figures 23, 28). To add the carboxy end of the AMV RT β subunit, since pAMVC has two *KpnI* sites (Figure 27), it was partially cleaved with *KpnI*, then completely cleaved with *ScaI*, and the 3 Kb fragment with the carboxy end of AMV RT was isolated and ligated to the 3.5 Kb *ScaI*-*KpnI* fragment of pAMVNMH- (Figure 29), forming pAMVBH- (Figures 23, 30).

Mutagenesis of the beta subunit to RNase H⁻. The RSV RT and AMV RT genes are related (GenBank sequences J02342, J02021 and J02343 for RSV-C; L10922, L10923, L10924 for AMV). These genes code for an identical sequence of amino acids over a short distance of the RNase H region. As described above in Example 1, during the cloning and mutagenesis of the RSV RT genes an RNase H⁻ derivative of the RSV RT β gene was made by site-directed mutagenesis. The oligonucleotide that was used (oligonucleotide #8) changed amino acid Asp450 to an Ala450 and introduced an *SstII* site (underlined).

Oligonucleotide #8

GGA CCC ACT GTC TTT ACC GCG GCC TCC TCA AGC ACC

The plasmid with the RNase H⁻ mutation in the RSV RT β gene is pAMP18BH- (Figures 2, 14). The 129 bp *BsrGI*-*BstEII* fragment from pAMP18BH- was cloned into the *BsrGI*-*BstEII* sites of pAMVNM, replacing the RNase H⁺ region in this plasmid and forming pAMVNMH⁻ (Figures 23, 29). The effect of this replacement was to change Asp450 to Ala450 in the AMV β gene without changing any other

amino acids. To convert the β subunit back to RNase H⁺, an oligonucleotide primer having the wildtype sequence may be used.

Mutagenesis and subcloning of the gene encoding the AMV RT α subunit.

To create a gene which codes for the α subunit of AMV RT, oligonucleotides #9 and #10 were used to mutagenize the amino end of the AMV RT gene from pAMVNM to introduce a translational stop codon where avian retroviral protease p15 normally cleaves the precursor polyprotein to make the α subunit:

Oligonucleotide #9 (SEQ ID NO:6):

5' CAU CAU CAU CAU CCC GGG TTA ATA CGC TTG GAA GGT GGC 3'

Oligonucleotide #10 (SEQ ID NO:7):

5' CUA CUA CUA CUA TCA TGA CTG TTG CGC TAC ATC TG 3'

PCR cycling conditions were 5 minutes at 94°C, followed by 8 cycles of 15 seconds at 55°C/2 minutes at 72°C/15 seconds at 94°C, and then 2 minutes at 72°C. The PCR reaction was treated with *DpnI* to destroy the target and the PCR product was cloned into pAMP18 by UDG cloning (Buchman, G.W., *et al.*, *Focus* 15:36 (1993)), forming plasmid pAMVA (Figures 24, 31). To make the RNaseH⁻ allele of the α subunit of AMV RT, the same procedure was followed using pAMVBH⁻ as a target, forming plasmid pAMVAH⁻ (Figures 24, 32).

Cloning the AMV RT α and β genes into pFastBac Dual. The α gene was excised from pAMVA with *SmaI* and *BspHI*, and subcloned into the *NcoI*-*PvuII* sites of pFastBac Dual (pD; Figure 33), creating plasmid pDAMVA (Figures 25, 34). An RNase H⁻ AMV RT α gene was similarly cloned from pAMVAH⁻, forming plasmid pDAMVAH⁻ (Figures 25, 35). In both plasmids, the AMV RT α gene was downstream from the baculovirus P10 promoter. The RNase H⁻ AMV RT β gene was excised from pAMVBH⁻ with *EcoRI* and cloned into the *EcoRI* site of pDAMVA. Clones were selected in which the *EcoRI* insert

was oriented such that the AMV RT β gene was downstream from the polyhedrin promoter, forming plasmid pDAMVABH- (Figures 25, 36). In this construct, the AMV RT α gene was RNase H⁺, but the β gene was RNase H⁻. The AMV RNase H⁻ RT β gene and polyhedrin promoter were excised from pDAMVABH- with *RsrII* and *NotI* and cloned into the *RsrII-NotI* sites of pSPORT1, forming plasmid pJAMVBH- (Figures 25, 37). The AMV RNase H⁻ RT β gene and polyhedrin promoter were excised from pJAMVBH- with *RsrII* and *NotI* and cloned into the *RsrII-NotI* sites of pDAMVAH-, forming plasmid pDAMVAH-BH- (Figures 25, 38).

The recombinant host cell comprising plasmid pDAMVABH-, *E. coli* DH10B(pDAMVABH-), was deposited on June 17, 1997, with the Collection, Agricultural Research Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois 61604 USA, as Deposit No. NRRL B-21790.

Using the deposited plasmid, one of ordinary skill in the art may easily produce, using standard genetic engineering techniques (such as those for site-directed mutagenesis described above), plasmids encoding various forms of the α and/or β subunits of AMV RT (*e.g.*, α RNase H⁺/ β RNase H⁺; α RNase H⁻/ β RNase H⁻; α RNase H⁺/ β RNase H⁻; and α RNase H⁻/ β RNase H⁺).

Transfection of insect cells for virus and AMV RT production. To prepare vectors for the transfection of insect cells, it was first necessary to insert the AMV RT gene constructs into a baculovirus genome. This insertion was accomplished using the site-specific transposon Tn7, as described for the insertion of the RSV RT gene constructs into bacmids in Example 1. Plasmid pDAMVABH- was transformed into DH10Bac cells and transposition of sequences encoding the AMV RT β gene (whose synthesis is directed by the baculovirus polyhedrin promoter) and the AMV RT α gene (whose synthesis is directed by the baculovirus P10 promoter) was followed by screening for loss of β -galactosidase activity following transposition to the site on the bacmid which codes for the β -galactosidase α peptide (Harris, R., and Polayes, D.A., *FOCUS* 19:6-8 (1997)).

Bacmid DNA was then prepared from transformants (10 ml cultures) by a slight modification of a standard miniprep procedure, as described in Example 1. About 1 µg of the bacmid DNA was used to transfect *Sf21* insect cells using the cationic lipid Cellfectin (Anderson, D., *et al.*, *FOCUS* 16:53 (1995)).

5 For expansion of primary virus, the supernatant was removed from the transfected cells about 72 hours after transfection and 1 ml was used to infect 35 ml of *Sf21* insect cells (about 1.2×10^5 cells/ml). After 72 hours, the culture was centrifuged (2,000 rpm; 10 min) and the supernatant was decanted and used as a secondary viral stock. The secondary virus stock was expanded similarly by
10 infecting 35 ml of *Sf21* cells with 0.1 ml of the secondary stock.

Virus stocks were then used to infect *Sf21* cells for the expression of AMV RT. In preliminary experiments, expression of RT activity by infected cells was found to be maximal about 72 hours after infection. For test expressions, 70 ml of *Sf21* cells were infected with 5 ml of the viral stock, and the cells were
15 harvested 72 hours after infection by centrifugation at 1,000 rpm for 5 minutes, resuspension in PBS (2.5% of the culture volume) and recentrifugation at 1,000 rpm for 5 min. Supernatants were removed and the cells were stored at -70°C until used. For larger scale production, 600 ml of cells in 2.8 liter Fernbach flasks were infected with 2 ml of viral stock, and the cells were harvested
20 72 hours after infection. AMV RT was then isolated as described for RSV RT in Example 2.

Example 5: Reverse Transcription with Retroviral RTs at Temperatures Above 55°C

25 Retroviral reverse transcriptases have historically been used to catalyze reverse transcription of mRNA at temperatures in the range of 37°C to 42°C (*see* technical literature of commercial suppliers of RTs such as LTI, Pharmacia, Perkin Elmer, Boehringer Mannheim and Amersham). There is a prevailing belief that at these temperatures mRNA secondary structure interferes with reverse

transcription (Gerard, G.F., *et al.*, *FOCUS* 11:60 (1989); Myers, T.W., and Gelfand, D.H., *Biochem.* 30:7661 (1991)) and the specificity of primer binding is reduced during gene-specific reverse transcription processes, such as RT-PCR, causing high background signal (Myers, T.W., and Gelfand, D.H., *Biochem.* 30:7661 (1991); Freeman, W.N., *et al.*, *BioTechniques* 20:782 (1996)). It is therefore desirable to carry out RNA reverse transcription at more elevated temperatures, *i.e.*, above 55°C, to help alleviate these problems.

As noted above, retroviral RTs are generally not used at temperatures above 37°C to 42°C to copy RNA because of the limited thermal stability of these mesophilic enzymes. In recent years, however, it has been reported that AMV RT can be used to perform RT-PCR of small amplicons (<500 bases) at 50°C, and to a limited extent at 55°C (Freeman, W.M., *et al.*, *BioTechniques* 20:782 (1996); Mallers, F., *et al.*, *BioTechniques* 18:678 (1995); Wang, R.F., *et al.*, *BioTechniques* 12:702 (1992)). Forms of M-MLV RT lacking RNase H activity, because of removal of the RNase H domain (Gerard, G.F., *et al.*, *FOCUS* 11:66 (1989) or because of point mutations in the RT gene (Gerard, G.F., *et al.*, *FOCUS* 14:91 (1992)), can also be used at 50°C, but not at 55°C, to catalyze cDNA synthesis.

Therefore, the thermal stability of RNase H⁻ RSV RT and its utility in higher temperature (*i.e.*, above 50°C) reverse transcription reactions for synthesis of large cDNAs was examined.

Methods

Enzymes and RNAs. SuperScript RT (SS RT), SuperScript II RT (SS II RT), and Moloney murine leukemia virus (M-MLV) RT were from LTI. AMV RT was from Seikagaku America, Inc., or was prepared as described above in Example 4. SS RT is an RNase H⁻ form of M-MLV RT in which RNase H activity has been eliminated by removing the RNase H domain of the RT polypeptide, resulting in an enzyme with a molecular weight of 57 KDa rather

than 78 KDa (Gerard, G.F., *et al.*, *FOCUS* 11:66 (1989); Kotewicz, M.L., *et al.* *Nuc. Acids Res.* 16:265 (1988)). SS II RT is an RNase H⁻ form of M-MLV RT in which RNase H activity has been eliminated by the introduction of three point mutations in the RNase H domain of M-MLV RT (Gerard, G.F., *et al.*, *FOCUS* 14:91 (1992)). Rous Associated Virus (RAV) RT was from Amersham. RSV RNase H⁻ and RSV RNase H⁺ RT were cloned, expressed and purified as described above in Examples 1 and 2. The RNAs used as templates were synthetic RNAs of 1.4, 2.4, 4.4 and 7.5 Kb, each with a 120 nucleotide poly(A) tail at the 3' end, obtained from LTI. Synthetic CAT mRNA was from LTI.

Model System for Determining Functional Thermal Stability of Reverse Transcriptases. A mixture of 1.4-, 2.4-, 4.4- and 7.5-Kb mRNAs was used to test the ability of various RTs to synthesize full-length cDNA copies at various temperatures. The cDNA products synthesized were labeled radioactively by the RT-catalyzed incorporation of a ³²P-labeled deoxyribonucleotide triphosphate precursor. The ³²P-labeled cDNA products were fractionated by alkaline agarose gel electrophoresis (Carmichael, G.G., and McMaster, G.K., *Meth. Enzymol.* 65:380 (1980)). The gel was dried and the size distribution of the cDNA products was established by autoradiography. Using the autoradiographic film as a template, the full length cDNA bands at 1.4, 2.4, 4.4 and 7.5 Kb were cut from the dried gel and counted in scintillant to establish the amount of each full length product synthesized.

cDNA synthesis reaction conditions. All cDNA synthesis reactions were carried out at the indicated temperatures for 30 or 50 minutes. All reaction mixtures were 20 µl and contained the following components unless specified otherwise: 50 mM Tris-HCl (pH 8.4 at 24°C), 75 mM KCl, 10 mM dithiothreitol, 1 mM each of [³²P]dCTP (300 cpm/pmole), dGTP, dTTP, and dATP, 25 µg/ml p(dT)₂₅₋₃₀, 12.5 µg/ml each of 1.4-, 2.4-, 4.4-, and 7.5-Kb mRNA, and 35 units of cloned rat RNase inhibitor. In addition, reaction mixtures contained the following:

SS II RT:	0.5 mM dNTPs (instead of 1 mM), 3 mM MgCl ₂ and 200 units of SS II RT
RSV H ⁻ RT:	7.5 mM MgCl ₂ and 21 units of RSV H ⁻ RT;
AMV RT:	50 mM KCl, 10 mM MgCl ₂ , 4 mM sodium pyrophosphate and 29 units of AMV RT
RSV H ⁺ RT:	50 mM KCl, 10 mM MgCl ₂ , 4 mM sodium pyrophosphate and 24 units of RSV H ⁺ RT
RAV RT:	50 mM KCl, 10 mM MgCl ₂ , 4 mM sodium pyrophosphate and 24 units of RAV RT

Reaction mixtures containing all components except enzyme were preincubated at the desired temperature for three minutes, and then RT was added to initiate cDNA synthesis.

Half Life Determinations. The half lives of RTs were determined by incubating individual tubes of RT at a desired temperature for appropriate lengths of time and stopping the incubation by placing the tube on ice. RSV RTs, RAV RT and AMV RT were incubated in 20 µl aliquots containing 50 mM Tris-HCl (pH 8.4), 75 mM KCl, 7.5 mM MgCl₂, 10 mM dithiothreitol, 50 µg/ml CAT mRNA, 25 µg/ml p(dT)₁₂₋₁₈, and 350-700 units/ml RT. Murine RTs (SS RT, SS II RT, M-MLV RT) were incubated in mixtures containing the same components except MgCl₂ was at 3 mM and enzyme was at 2,500 units/ml. An aliquot from each tube (5 µl for avian RTs and 1 µl for murine RTs) was assayed for RT activity in a unit assay reaction mixture to determine residual RT activity.

Unit Assays. Unit assay reaction mixtures (50 µl) contained 50 mM Tris-HCl (pH 8.4), 40 mM KCl, 6 mM MgCl₂, 10 mM dithiothreitol, 500 µM [³H]dTTP (30 cpm/pmole), 0.5 mM poly(A), and 0.5 mM (dT)₁₂₋₁₈. Reaction mixtures were incubated at 37°C for 10 minutes and labeled products were acid precipitated on GF/C glass filters that were counted in a scintillation counter.

Results and Discussion

With a few exceptions, the half lives in the presence of a template primer at 45°C, 50°C, 55°C and 60°C were determined for RSV H⁺ RT, RAV RT, AMV RT, RSV H⁻ RT, M-MLV H⁺ RT, SS RT, and SS II RT. The results are shown in Figure 39 and Table 3.

TABLE 3. HALF LIVES OF REVERSE TRANSCRIPTASES ¹				
ENZYME	HALF LIFE (MINUTES) AT:			
	45°C	50°C	55°C	60°C
RSV H ⁻ RT	440	138	5	0.75
RSV H ⁺ RT	ND ²	30	ND ²	ND ³
RAV RT	159	37	3.8	ND ³
AMV RT	96	16	1.3	ND ³
SuperScript II RT	105	7	ND ³	ND ³
SuperScript RT	120	3	ND ³	ND ³
M-MLV RT	65	2.5	ND ³	ND ³

¹Half lives were determined from the data shown in Figure 39.

²ND: not determined.

³ND: not determined, because half life in this reaction mixture was too short to be determined accurately.

The results shown in Figure 39 and Table 3 clearly demonstrate that RNase H⁺ RTs (RSV, RAV and AMV) are much more thermostable than M-MLV RT, and have reasonable half lives at 50°C. Furthermore, mutating these RTs to produce their corresponding RNase H⁻ forms further increases their half lives. Most dramatically, RNase H⁻ RSV RT had a much longer half life than any other retroviral RT at 45°C, 50°C and 55°C. Thus, introduction of a single amino acid change into the RNase H domain of each subunit of RSV RT increases its half life at 50°C by nearly five-fold.

The impact of this increased thermal stability on cDNA synthesis at temperatures above 50°C was found to be dramatic. Figure 40 shows the results

of a comparison of the performance of these RTs at 45°C, 50°C, 55°C and 60°C in copying mRNA of 1.4 to 7.5 Kb in length. With the exception of RSV H⁻ RT, none of the RTs were found to produce substantial product longer than 2.4 Kb in length at 55°C or 60°C. RSV H⁻ RT, in contrast, continued to make full-length 7.5-Kb cDNA at 55°C, and 4.4-Kb cDNA at 60°C. Figures 41 and 42 show a more detailed comparison of the two RTs that performed best in Figure 40 (*i.e.*, SS II RT and RSV H⁻ RT). At temperatures above 55°C, RSV H⁻ RT was found to continue to synthesize cDNAs of all lengths, while SS II RT produced only low levels of cDNA greater than 1 Kb in length.

Taken together, these results demonstrate that RNase H⁻ RSV RT is much more thermoactive than any other retroviral RT available commercially, and can be used to synthesize longer cDNAs (up to 4 Kb long) at 60°C. This enhanced thermoactivity of RNase H⁻ RSV RT is due to an increased thermal stability, relative to RNase H⁺ RT, at 50°C to 60°C, which makes RSV H⁻ RT an ideal enzyme for use in the reverse transcription of mRNA at 50°C to 60°C.

Example 6: Reverse Transcription with Avian RTs at Elevated Temperatures

In Example 5 (Table 3 and Figure 39), the half lives of various RTs were presented. In particular, half lives were reported for cloned RSV RT in which the RNase H domain of each subunit was mutated to eliminate RNase H activity (Example 1; Asp450 → Ala in both the α subunit and the β subunit).

To further examine the effects of these mutations on RT half life, constructs were produced as described above, in which only one of the two subunits were mutated at one time, such that various combinations of mutants were formed (*e.g.*, α RNase H⁻/β RNase H⁺ and α RNase H⁺/β RNase H⁻). The half lives of these RSV RTs, as well as cloned AMV α RNase H⁻/β RNase H⁺ RT, were determined as described above in the Methods section of Example 5.

As shown in Table 4, when the α subunit in RSV or AMV RT was mutated, leaving the β subunit wild type intact, the resulting RT demonstrated greater thermal stability than that observed for other avian RTs. These mutant RTs were also examined for their functional thermal stability using the model system described in the Methods section of Example 5. For each enzyme, the increased thermal stability was found to correlate with improved functional performance -- for example, the α RNase H/ β RNase H⁺ avian RTs, which demonstrated the highest thermal stability at 55°C (Table 4), also demonstrated the highest functional activity at various elevated temperatures (Table 5).

Table 4. Half Lives of RSV and AMV Reverse Transcriptases	
Enzyme	Half Life (Minutes) at 55°C
RSV α H β H RT	5
RSV α H β H ⁺ RT	7
RSV α H ⁺ β H RT	2
RSV α H ⁺ β H ⁺ RT	1.9
AMV α H β H ⁺ RT	6
Native AMV RT	1.3
Native RAV RT	3.8

Table 5. Functional Activities of RSV RTs at Elevated Temperatures.					
Enzyme	Temperature, °C	Amount of Full-Length Product Produced (pMoles)			
		1.4 Kb	2.4 Kb	4.4 Kb	7.4 Kb
RSV α H/ β H ⁻ RT	45.0	132.9	80.5	56.7	28.1
	55.0	115.4	70.7	40.8	12.5
	57.5	81.9	43.2	17.2	3.1
	60.0	7.0	1.8	0	0
	62.5	0	0	0	0
RSV α H/ β H ⁺ RT	45.0	145.2	85.3	57.9	31.8
	55.0	161.3	83.0	53.3	21.5
	57.5	140.1	77.7	41.1	11.6
	60.0	67.6	30.0	7.5	0.1
	62.5	4.1	0.8	0	0

Example 7: Alternative Methods of Generating Avian Reverse Transcriptases and Characterization of their Properties

As noted above in the Related Art section, three prototypical forms of retroviral RT have been studied thoroughly -- M-MLV RT, HIV RT, and ASLV RT (which includes RSV and AMV RT). While each of these retroviral RTs exist as heterodimers of an α and a β subunit, there have been no reports heretofore of the simultaneous expression of cloned ASLV RT α and β genes resulting in the formation of heterodimeric $\alpha\beta$ RT.

Examples 1-4 above described the cloning, expression and purification of $\alpha\beta$ forms of RSV and AMV RT that copy mRNA efficiently. Formation of $\alpha\beta$ RT was achieved in baculovirus-infected insect cells by co-expression of genes for α

and β from a dual promoter vector. The studies presented in this Example were designed to generate RSV $\alpha\beta$ RT by a variety of other methods that have been used to successfully clone and express HIV p66/p51 RT. In addition, as described below, the individual subunits of RSV RT, including β p4, β and α , have now been cloned, expressed, and purified, and the abilities of these subunits to copy mRNA has now been characterized.

MATERIALS AND METHODS

General Methods

Mutations and plasmid constructions were conducted using standard molecular biology methods (see e.g., Sambrook, J. *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989)), modified as described below. Plasmid preparation, PCR, gel electrophoresis, DNA fragment isolation and cloning, insect cell culture and baculovirus production were all performed as described for RSV RT cloning and expression in Example 1.

Cloning and Expression of RSV RT in E. coli

A number of approaches were tried to generate RSV $\alpha\beta$ RT from RSV RT β p4 in *E. coli*.

PCR of NdeI-XbaI fragment. The amino end of the RSV RT β p4 gene was mutagenized to introduce an *NdeI* site by PCR (1 cycle 94° C for 5 min.; 15 cycles 94° C, 10 sec, 55° C, 15 sec, 72° C, 15 sec.; 1 cycle 72° C, 5 min) of pJD100 (Figure 7) with the following oligonucleotides:

Oligonucleotide #11 (SEQ ID NO:17):

5'- ATT ATT CAT ATG ACT GTT GCG CTA CAT CTG GC-3'

Oligonucleotide #12 (SEQ ID NO:18):

5'- TAC GAT CTC TCT CCA GGC CAT TTT C-3'

The *NdeI* site in oligonucleotide #11 above appears in bold and italicized print, while the bases that are underlined in oligonucleotides #11 and #12 were derived from authentic RT gene sequences. The PCR product with these two oligos contained an *NdeI* site at the beginning of the gene, and retained the *XbaI* site which is present within the RT gene. The PCR product was cloned into pUC18 between the *NdeI* and *XbaI* sites, forming pUC18#3.

Site-directed mutagenesis to introduce a SmaI site to clone p15. A 3 Kb *PstI* fragment from pJD100 containing the entire RSV RT gene was cloned into M13mp19. Clone RF-*SmaI* was then produced by introducing a *SmaI* site at the carboxy end of the RSV RT β p4 gene by site-directed mutagenesis (see Example 1), using the following mutagenic oligonucleotide:

Oligonucleotide #13 (SEQ ID NO:19):

5'- ACT CGA GCA GCC **CGG** GAA CCT TTG -3'

Reconstruction of the RT gene. The 2.8 kb *SmaI-PstI* fragment from RF-*SmaI* was cloned into pUC18 at the *SmaI/PstI* sites. The clone was designated as pUC18-*PstI-SmaI*. The *NdeI-XbaI* fragment from pUC18 #3 was introduced into pUC18-*PstI-SmaI* to regenerate the entire RSV RT β p4 gene with an *NdeI* site at the initiation codon. The clone was designated as pUC18-RT.

Cloning entire RT gene in pRE2. The *NdeI-XbaI* fragment from pUC18 #3 was cloned into expression vector pRE2 to generate pRE2-Nde-Xba. pRE2 contains an inducible lambda pL promoter. The rest of the RSV RT β p4 gene was subcloned as an *NdeI-SstI* fragment from pUC18-*PstI-SmaI* into pRE2-Nde-Xba digested with *XbaI* and *SstI*, generating pRE2-RT.

Cloning of p15 into pRE2-RT. RSV RT is processed from RSV RT β p4 to the $\alpha\beta$ heterodimer by the RSV p15 protease. In order to make authentic $\alpha\beta$,

the p15 protease gene was cloned and expressed with the RSV RT β p4 gene as follows. The RSV p15 protease gene was generated by PCR using pJD100 as target and the protocol described above. The oligonucleotides used for PCR were as follows:

Oligonucleotide #14 (SEQ ID NO:20):

5'-AT TAC **CCG** GGA GG A TAT CAT ATG TTA GCG ATG ACA ATG
GAA CAT AAA G -3'

Oligonucleotide #15 (SEQ ID NO:21):

5'- A TAT **GTC GAC** TCA CAG TGG CCC TCC CTA TAA ATT TG-3'

In oligonucleotides #14 and #15 above, the restriction sites (*Sma*I, *Nde*I and *Sal*I) are indicated in bold letters while the region of bases underlined is the ribosome binding site. PCR using these oligonucleotides generated a ~ 450 bp fragment, which was digested with *Sma*I and *Sal*I and cloned into pUC19. The clone was designated as pUC19-p15. The p15 gene was introduced into pRE2-RT by subcloning the 450 bp *Sma*I-*Sal*I fragment at the *Sma*I/*Sal*I sites. The final plasmid was designated pRE2-RT.p15.

Expression of RT from pRE2-RT and pRE2-RT.p15. *E. coli* CJ374 containing either pRE2-RT or pRE2-RT.p15 was grown at 30°C in EG broth in the presence of ampicillin (100 µg/ml) and chloramphenicol (30 µg/ml) to an A590 of 0.5. Half of the culture was induced at 42°C for 45 min., and then outgrown at 30°C for 2 hr. The other half was grown at 30°C as an uninduced control. None of the cultures produced any visible induced protein upon examination by SDS-PAGE. None of the cell extracts displayed any RT activity.

Recloning of RT.p15 in pRE1. The lack of RT expression in the above constructs suggested that it was possible that (i) a mutation had been introduced into the RT gene during PCR to render it inactive, or (ii) during cloning a

mutation in the lambda pL promoter arose, since RT is thought to be toxic to *E. coli*. Thus, the entire RT.p15 gene was recloned into pRE1, which was the same as pRE2 except the multiple cloning site was in an opposite orientation, as a *NdeI-SaII* fragment and the construct was introduced into *E. coli* CJ374. In addition, a 2269 bp *HpaI-KpnI* fragment of the resulting clone was replaced by the same *HpaI-KpnI* fragment from pJD100. This replacement left only about a 200 bp amino terminal region derived from PCR. Moreover, this region (*NdeI* site to *HpaI* site) was sequenced to confirm that there was no mutation due to PCR. The plasmids were designated as pRE1-RT.15. This plasmid was also introduced into BL21, a protease-deficient *E. coli* strain. The p15 gene was also deleted from pRE1-RT.15 by digesting the plasmid with *XhoI* and *SaII* and recircularizing the plasmid. The resulting plasmid was designated as pRE1-RT.

Expression of RT in E. coli CJ374 and BL21 harboring pRE1-RT and pRE1-RT.15. The cultures were grown as described above. The soluble cell extract was assayed for RT activity. Although the activity was extremely low, it was clear that the RT activity in the induced cell extract was 10-times higher than that in the uninduced cell extract. The level of activity was similar in both CJ374 and BL21. The levels of expression of RT from pRE1-RT and pRE1-RT.15 were similar.

Cloning of RT gene under a tac promoter. Since RSV RT β p4 was not expressed well under a lambda pL promoter, expression was attempted under the control of a different promoter. The RSV RT β p4 gene with and without the p15 gene was cloned under a *tac* promoter. For cloning the RT gene, pRE1-RT was digested with *NsiI*, blunt-ended with T4 DNA polymerase and finally, digested with *XhoI*. The RT fragment was purified from an agarose gel. For cloning the RT.15 gene, pRE1-RT.15 was digested with *NsiI*, blunt-ended with T4 DNA polymerase and finally, digested with *SaII*. The RT.15 gene combination fragment was purified from an agarose gel. The vector pTrc99A (Pharmacia) was digested with *NcoI*, blunt-ended with Klenow fragment and finally, digested with *SaII*. The

large vector fragment was purified and ligated with either purified RT or RT.15 fragment. The resulting constructs were introduced into *E. coli* DH10B. Clones with correct inserts were saved. The clones were designated as pTrcRT and pTrcRT.15.

5 *Expression of RT under a tac promoter.* *E. coli* cells harboring pTrcRT or pTrcRT.15 were grown at 37°C in a buffered-rich medium to an A590 to 0.1 or 0.6 before addition of IPTG (1mM) for induction of RSV RT β p4 expression. The cells were collected 2 hr after induction. The uninduced cultures were grown similarly without addition of any inducer. The RT enzyme activity in soluble cell
10 extract was equivalent to that obtained from constructs with the lambda pL promoter.

SDS polyacrylamide gel electrophoresis of the expressed proteins. RSV RT is composed of two subunits of molecular mass 94 kD (β) and 62 kD (α). The α subunit is a proteolytic fragment derived from the β subunit. The proteolysis is accomplished by RSV p15 protease *in vivo*. However, when
15 induced *E. coli* extracts bearing the plasmids described were examined, two induced proteins of molecular mass 75 kD and 62 kD were detected. Both proteins were found to be mainly in the insoluble fraction of *E. coli* extracts. The presence of a 75 kD and not the expected 94 kD protein following expression in
20 *E. coli* suggested that (i) there was a mutation near the carboxy end causing premature translation termination, (ii) there was proteolysis in the *E. coli* cytosol, or (iii) there was internal translation initiation of RT. When the carboxy terminal region of the RT gene was sequenced, no mutation was found that might have caused premature translation termination. To test whether there was an internal
25 translation start, the RT gene was cloned as a *HpaI-XhoI* fragment in pTrc99 digested with *SmaI* and *SalI*. No induced protein could be detected suggesting that there was no internal start to generate either a 75 kD or 62 kD protein.

Expression of RT in a variety of E. coli hosts. As described above, expression of RSV RT was examined in *E. coli* DH10B, CJ374 and BL21. None

of these *E. coli* hosts produced $\alpha\beta$ RT, and the level of RT activity in each host was extremely low. Since M-MLV RT expresses well in *E. coli* N4830 and HIV RT expresses well in *E. coli* RR1, the levels of expression of RSV RT in these two hosts were examined. Neither of these hosts was found to express active RT any better than the other hosts tested, nor did they produce any full length 94 kD protein.

Expression of RT as a fusion protein. It is now well established that some proteins that do not express well in *E. coli* are better expressed as a translation fusion, in which the protein from a well expressed gene forms the amino end of the fusion protein. The gene of one such fusion partner, thioredoxin, is present in a vector called pTrxfus (Genetics Institute; Cambridge, MA). The level of expression of RSV RT β p4 fused with thioredoxin was therefore examined.

In the course of expressing RSV RT in a baculovirus system, the RSV RT gene fragment was cloned as a *SmaI-XhoI* fragment in pBacPAK9 to construct pBacPak-RT (see below). To make the thioredoxin fusion, pBacPak-RT was digested with *XmaI* (same recognition as *SmaI*) and *PstI* and the RT fragment was purified on an agarose gel. The vector pTrxfus was digested with *XmaI* and *PstI* and the large vector fragment was purified. The purified fragments were ligated and *E. coli* CJ374 and GI724 or GI698 (Genetic Institute, MA) were transformed with the ligated material. Clones with the correct insert were saved. When cultures were induced and the cellular extracts were assayed for enzyme activity, no RT activity was detected. Both induced cultures, however, produced a large amount of insoluble fusion protein of the expected size as judged by SDS-PAGE of the extracts. Other fusions were also tested. The RSV RT β gene was fused to the GST (glutathione S transferase) gene, and the RSV RT α gene was fused to the lambda CRO protein gene. Expression of either fusion from the *trc* promoter in *E. coli* strain DH10B resulted in a large amount of insoluble protein of the appropriate molecular weight, but little RT activity was detected in extracts of induced cells. An expression vector with both fusions (GST-beta and CRO-

alpha) was constructed in which the fusions were co-expressed by induction of the *trc* promoter. Co-expression of both fusions in *E. coli* strain DH10B resulted in large amounts of insoluble protein of the appropriate molecular weights, but little RT activity.

5 *Cloning and Expression Separately of Genes Encoding RSV RT Subunits α , β , and $\beta p4$ in a Baculovirus System*

Cloning of RSV RT $\beta p4$ in a baculovirus system

To clone the RSV RT $\beta p4$ gene in the baculovirus transfer vector, pBacPAK9 (Clontech), a fragment of the RSV RT gene was generated by PCR. To facilitate this PCR, the following oligonucleotide was designed with a *Bam*HI site (bold) and the ATG initiation codon at the beginning of the RSV $\beta p4$ gene:

Oligonucleotide #16 (SEQ ID NO:22):

5'-TAT TAG **GAT CCC ATG** ACT GTT GCG CTA CAT CTG GC-3'

Oligonucleotides #12 and #16 (SEQ ID NOs: 18 and 22, respectively) were used for PCR using pJD100 as template. The PCR product was digested with *Bam*HI and *Xba*I, and then ligated to pBacPAK9 digested with *Bam*HI and *Xba*I. One of the clones, pBP-RT(PCR) (Figure 43), was used for further cloning. To reconstitute the entire RT gene, the small *Hpa*I-*Xho*I fragment of pBP-RT(PCR) was replaced with the 2500 bp *Hpa*I-*Xho*I fragment of pRE1-RT.15. The reconstituted plasmid was designated as pBP-RT(ATG) (Figure 44).

Cloning of p15 in pBacPAK-RT(ATG). Placing the RSV p15 protease gene in pBP-RT(ATG) was achieved by a three-step cloning. First, a p15 gene fragment was subcloned from pUC19-p15 (see above) into pSport1 (LTI) as a *Kpn*I-*Sal*I fragment to generate pSport-p15. Second, the p15 fragment was subcloned from pSport-p15 as a *Nof*I-*Sma*I fragment into a baculovirus transfer vector, pAcUW43, to generate pAcUW43-p15. This cloning was done to introduce the p15 gene under a p10 promoter. Finally, the p15 gene including the

p10 promoter was subcloned from pAcUW43-p15 as a *XhoI*-*NoI* fragment into pBP-RT (ATG) to generate pBP-RT15(ATG) (Figure 45).

Expressing RSV RT β p4 in insect cells. Insect cells (SF9) were transfected with a mixture of pBP-RT15(ATG) and linearized baculovirus vector DNA (BaculoGold, Pharmingen). In this system, recombination between the pBP-RT15(ATG) plasmid and the baculoviral DNA results in recombinant viral genomes with the RSV RT β p4 gene downstream of the polyhedrin promoter. Recombinant virus was isolated from the supernatant by standard techniques and viral stocks were made, one of which was chosen for further study. Insect cells were infected with pure viral stock and infected cells were harvested at various times after infection. RT activity was easily detectable in the infected cells.

Generation of an RNase H- RSV RT. The generation of a mutation in the RNase H region of the RSV RT β p4 gene is described in detail above in Example 1. To introduce the mutation into pBP-RT(ATG), the *HpaI*-*KpnI* fragment of pBP-RT(ATG) was replaced with the *HpaI*-*KpnI* fragment of M13RTH- (Figure 13). The newly constructed plasmid was designated as pBP-RT(H-) ATG. Insect Cells (SF9) were transfected with a mixture of pBP-RT(H-) and linearized baculovirus vector DNA (BaculoGold, Pharmingen). Recombination between the pBP-RT(H-) plasmid and the baculoviral DNA results in recombinant viral genomes with the RSV RT β H-P4 gene downstream of the polyhedrin promoter. Recombinant virus was isolated from the supernatant by standard techniques. Insect cells were infected with viral stock and infected cells were harvested at various times after infection. RT activity was easily detectable in the infected cells.

Attaching a histidine tag to RSV RT β p4. pBP-RT(H-) was cleaved with *Bam*HI and *Xba*I and the 0.9 kb fragment with the amino end of the RSV RT β p4 gene was cloned into pFastBacHT at the *Bam*HI-*Xba*I sites, creating (by translational fusion) a β p4 partial construct with a histidine tag at the amino end (pFBHT β P4). The 2 kb *Xba*I-*Xho*I fragment with the carboxy end of RSV RT β H-p4 from pBP-RT(H-) was inserted into the *Xba*I-*Xho*I sites of pFBHT β P4,

creating pFBH-TβP4. pFBH-TβP4 was transformed into *E. coli* DH10Bac cells and the His-tagged RSV RT βH-p4 gene was transposed to bacmid. Bacmid DNA was isolated and transfected into Sf9 insect cells. Viral preparations from infected cell cultures were used to infect SF21 cells, and His-tagged RSV RT βH-p4 from infected cells was isolated and characterized.

Transfection of insect cells. The transfection of Sf9 cells was done using Baculogold virus (Pharmingen, CA) and pBP-RT(H-)ATG to generate recombinant virus. Ten wells on two 6-well (60 mm) tissue culture plates were seeded 1×10^6 Sf9 cells (LTI). The cells were allowed to attach for 30 min at 27°C. While the cells were attaching, ten tubes each containing 200 μl of Sf-900II SFM medium (LTI) were set up. In tube 1, 500 ng of Baculogold and 2 μg of pBP-RT(H-)ATG were added. In tube 2, 250 ng of Baculogold and 1 μg of pBP-RT(H-)ATG; and tube 3, 125 ng of Baculogold and 500 ng of pBP-RT(H-)ATG were added. Tubes 6 through 9 contained 36 μl of lipofectin (LTI). Contents of tube 1, tube 2, tube 3, tube 4 and tube 5 were transferred to tubes 6, 7, 8, 9 and 10, respectively. Into each of these tubes, 2 ml of Sf-900 II SFM were added. The culture medium from each well was removed and 2 ml of the DNA/lipofectin mix was dispensed in two wells, 1 ml each. Thus, the wells containing the mixture of tubes 1 and 6, 2 and 7, and 3 and 8 contained DNA mixtures at different amounts; the wells containing the mixtures of tubes 4 and 9 were controls containing lipofectin but no DNA; and the wells containing the mixtures of tubes 5 and 10 were controls containing neither DNA nor lipofectin. The plates were incubated at 27°C for 4 hr, the medium was removed and replaced with 4 ml fresh Sf-900 II SFM, and the plates were incubated for an additional 72 hrs at 27°C. The phage supernatants from the DNA containing wells were removed and marked as primary phage stocks. A second infection was done by infecting 1×10^6 Sf9 cells with 1 ml of the primary phage stock to amplify the recombinant phage. The plates were incubated for 48 hrs at 27°C and the phage stocks were collected.

Injection of insect larvae. Trichoplusia ni larvae were injected with recombinant virus bearing pBP-RT(H-)ATG and the larvae were harvested as described (Medin, J.A., *etal.*, *Methods in Molecular Biology* 39:26 (1995)).

Expression of RSV RT α H- in a baculovirus system

pDA (Figure 19) was transformed into *E. coli* DH10Bac cells, the RT α H-gene was transposed to bacmid, and the bacmid DNA was purified and transfected into SF21 insect cells. Viral preparations from infected cell cultures were used to infect SF21 cells, and RSV RT α H- from infected cells was isolated and characterized.

Expression of RSV RT β H-His in a baculovirus system

pDABH-His (Figure 21) was cleaved with *Rsr*II and *Pst*I, and the 2.6 kb fragment with the β H-His gene was cloned into the *Rsr*II-*Pst*I sites of pFastBac. pFBBH-His was transformed into *E. coli* DH10Bac cells, the RT β H-His gene was transposed to bacmid, and the bacmid DNA was purified and transfected into SF21 insect cells. Viral preparations from infected cell cultures were used to infect SF21 cells, and RSV RT β H-His from infected cells was isolated and characterized.

Cloning and Expression of Genes Encoding RSV $\alpha\beta$ RT in which the Polymerase Active Site is Mutated

Generation of RSV RTs mutated in the polymerase domain. Alignment of the RSV RT peptide sequence with sequences from HIV RT, M-MLV RT, and sequences of other RT genes revealed the probable location of one of the catalytic residues in the RSV RT polymerase domain, D110 (aspartic acid residue at position number 110). According to the literature, mutation of the corresponding amino acid in the larger chain of HIV RT from D (aspartate) to E (glutamic acid)

resulted in nearly total loss of polymerase activity. Single strand DNA was isolated from pJB-His by infection of *E. coli* DH5 α F'IQ/pJB-His cells with M13KO7, and this DNA was mutated by site-directed mutagenesis (see Example 1 for detailed protocol) with the following oligonucleotide :

5 Oligonucleotide #17 (SEQ ID NO:23):
5'- GCAATCCTT**GAGCTCT**AAGACCATCAGGG 3'

This oligonucleotide induces the mutation of the aspartate residue at position #110 or the RSV RT catalytic domain to glutamate (D110E) and adds an *Sst*I site (bold), forming plasmid pJBD110E-His (Figure 48). This mutated site was introduced into the RSV RT α gene by inserting the 460 bp *Nhe*I-*Eco*47III fragment from pJBD110E-His into the 6.5 kb fragment of *Nhe*I-*Eco*47III cleaved pDA, forming pDAD110E (Figure 49). The D110E mutation was also introduced into the β gene by inserting the 460 bp *Nhe*I-*Eco*47III fragment into *Nhe*I-*Eco*47III cleaved pFBBH-His (Figure 46), forming pFBBD110E-His (Figure 50). The 2.6 kb RSV RT β D110E gene was cloned from pFBBD110E-His into pDABHis (Figure 51) as a 2.6 kb *Rsr*II-*Eco*RI fragment, replacing the RSV RT β -His gene, forming pDABD110EHis (Figure 52). pDABHis was cleaved with *Xho*I + *Pvu*I, and the 4.6kb fragment with the β gene was joined to the 4.9 kb pDAD110E *Xho*I-*Pvu*I fragment, forming pDAD110EBHis (Figure 53). The 4.6 kb pDABD110EHis *Xho*I-*Pvu*I fragment with the β D110E gene was joined to the 4.9 kb pDAD110E *Xho*I-*Pvu*I fragment, forming pDAD110EBD110E (Figure 54) (which has a his tag despite its truncated name). pDAD110EBHis, pDABD110EHis, and pDAD110EBD110E were transformed into *E. coli* DH10B-Bac, the RT genes were transposed to the bacmid, and the bacmid DNA was purified and transfected into SF21 insect cells. Viral preparations from infected cell cultures were used to infect SF21 cells, and RSV RT from infected cells was isolated and characterized.

The three mutant plasmids are summarized in the following table, where "w.t." refers to the wild type amino acid at the 110 position (D, aspartic acid) and D110E refers to the mutation at the 110 position (D to E, glutamic acid):

<u>Plasmid</u>	<u>α</u>	<u>β</u>
pDABHis	w.t.	w.t.
pDABD110EHis	w.t.	D110E
pDAD110EBHis	D110E	w.t.
pDAD110EB110E	D110E	D110E

Cloning and Expression of RSV β p4 RT in Yeast

Cloning of the RSV β p4 RT gene. The pHIL-D2 vector available from InVitrogen (CA) was digested with EcoRI, blunt ended with Klenow fragment and treated with alkaline phosphatase. pRE1-RT.15 was digested with *Nde*I and blunt ended with Klenow fragment. The *Nde*I digest generated the entire RSV RT gene without the p15 protease gene. The fragments were ligated and *E. coli* DH10B was transformed with ligation mixture. The correct clones were selected for proper insert and orientation. Two of the 8 clones tested had the RT gene fragment in the correct orientation. One of the clones, pHILD2-RT, was used for further experimentation. To introduce the RNase H- domain in this plasmid, pHILD2-RT was digested with *Bam*HI plus *Xho*I and the wild type fragment was replaced with *Bam*HI-*Xho*I fragment from pBacPAK-RT(H-)ATG. The final clone was screened with *Sst*II. The clone was designated as pHILD2-RT(H-).

Transformation of Pichia pastoris. *Pichia pastoris* GS115 (InVitrogen) was used for transformation according to the protocol recommended by InVitrogen. The plasmids, pHILD2-RT and pHILD2RT(H-), were digested with *Not*I, phenol-chloroform extracted and ethanol precipitated before transformation. The transformation yielded 20 clones for wild type RSV RT and 12 clones for

RSV H- RT in the regeneration plates. These clones were screened for their growth in methanol containing plates. Two putative clones were selected from initial 20 clones (wild type RT). One of these two, H1, was completely incapable of growing in methanol, and the other, H2, was capable of growing very slowly in methanol. For the RSV H- RT clones, three clones were selected out of 12 screened. One of the clones, H-3, grew very slowly in methanol. The others, H-4 and H-5, grew moderately in methanol. Clones H1 and H-3 were chosen for expression studies.

RSV $\beta p4$ RT expression in Pichia pastoris. Clones H1 and H-3 were grown and induced essentially as described in the user's manual provided by the manufacturer (InVitrogen). As a control, GS115 containing a β -galactosidase gene (InVitrogen) was grown and induced side by side. While no RT activity could be detected in the GS115/ β -gal cells, an appreciable level of activity was detected in both H1 (wild type RT) and H-3 (H- RT) cells. In addition, the activity increased with increased time of induction.

Isolation of RSV $\beta\beta$, α and $\beta p4\beta p4$

RSV $\beta\beta$ RT isolation. RSV $\beta\beta$ RT was purified as described in Example 2 for RSV $\alpha\beta$ RT with the following exceptions. RSV $\beta\beta$ RT eluted from the AF-Heparin-650M column from 55-62% Buffer T, and from the Mono S HR 5/5 column from 58-62% Buffer J. RSV $\beta\beta$ RT was 90% homogeneous as judged by SDS-PAGE.

RSV α RT isolation. RSV α RT was purified as described in Example 2 for RSV $\alpha\beta$ RT with the following exceptions. The Chelating Sepharose Fast Flow column was equilibrated with 100% Buffer A and washed with 100% Buffer A after the clarified crude extract was passed over the column. RSV α RT was eluted with a 10-column volume linear gradient of 100% Buffer A to 75% Buffer A + 25% Buffer B. The peak fractions of RT activity from the Chelating Sepharose Fast Flow column (10-12% Buffer B) were pooled, dithiothreitol and EDTA were added to the pool to achieve final concentrations of 1 mM and 0.1

mM, respectively, and the pool was dialyzed overnight against 95% Buffer S + 5% Buffer T. The dialyzed pool was loaded on a 22 ml AF-Heparin-650M column equilibrated in Buffer S. After a wash with 9 column volumes of 95% Buffer S + 5% Buffer T, the column was eluted with a 9 column volume linear gradient of 95% Buffer S + 5% Buffer T to 60% Buffer S + 40% Buffer T. The peak fractions of RT activity (11-20% Buffer J) were pooled and dialyzed for 3 to 4 hours against 97.5% Buffer H and 2.5% Buffer J. The dialyzed pool was loaded onto a 3.5 ml phosphocellulose column (Whatman) equilibrated in 100% Buffer H. After a wash with 12 column volumes of 97.5% Buffer H + 2.5 Buffer J, the column was eluted with a 14 column volume linear gradient of 97.5% Buffer H + 2.5% Buffer J to 60% Buffer H + 40% Buffer J. The peak fractions of RT activity (12-20% Buffer J) were pooled and dialyzed overnight against 99% Buffer S + 1% Buffer T. The dialyzed pool was loaded onto a Mono S HR 5/5 column equilibrated in Buffer S. After a wash with 10 column volumes of 100% Buffer S, the column was eluted with a 20 column volume linear gradient of 100% Buffer S to 75% Buffer S + 25% Buffer T. The RT peak fractions (15-17% Buffer T) were pooled, dialyzed against Storage Buffer overnight, and stored at -20°C. RSV α RT was found to be 80% homogeneous as judged by SDS-PAGE.

RSV $\beta p4\beta p4$ RT isolation. RSV $\beta p4\beta p4$ RT was purified as described in Example 2 for RSV $\alpha\beta$ with the following exceptions. The pooled RT fractions from the Chelating Sepharose Fast Flow column were dialyzed overnight against 90% Buffer H + 10% Buffer J. The RSV $\beta p4\beta p4$ RT precipitated in this buffer. The RT was recovered by centrifugation, dissolved in Storage Buffer containing 0.5 M KCl, and stored at -20°C. RSV $\beta p4\beta p4$ RT was found to be > 95% homogeneous as judged by SDS-PAGE.

Estimation of the Amounts of α , $\alpha\beta$ and $\beta\beta$ RSV RT in Baculovirus-Infected Insect Cells

Chromatography methods described in Example 2 can be used to separate and isolate any α , $\alpha\beta$ and $\beta\beta$ RSV RT present in crude extracts from virus-infected insect cells. RSV RT α is separable from the other two enzymes forms by chromatography on a Chelating Sepharose Fast Flow column, where α either does not bind or elutes at $< 30\text{mM}$ imidazole, and $\alpha\beta$ and $\beta\beta$ elute together at $> 50\text{ mM}$ imidazole. This is possible by virtue of the His₆ tag present on β , but not on α . RSV $\alpha\beta$ and $\beta\beta$ RT are separable subsequently by chromatography on Heparin-650M ($\alpha\beta$ elutes at 0.45M KCl and $\beta\beta$ elutes at 0.58M KCl). Reverse transcriptase is quantitated by assay with poly(C)•oligo(dG) (Gerard, G.F., *et al.*, *Biochemistry* 13: 1632 (1974)). The specific activities of RSV RT α , $\alpha\beta$ and $\beta\beta$ with poly(C)•oligo(dG) are 140,000, 90,000 and 6,000 units/mg of protein, respectively.

Cloning, Expression, Purification and Use of RSV Viral Protease p15

RSV viral protease was cloned and expressed as a linked dimer in *E. coli* and purified from inclusion bodies as described (Bizub, D., *et al.*, *J. Biol. Chem.* 266: 4951 (1991)).

Reaction mixtures ($25\mu\text{l}$) used to digest RSV βp4 RT with RSV protease contained 100 mM NaPO_4 (pH 6.0 to 7.0), 1 mM 2-mercaptoethanol, 0.01% (W/V) Triton X-100, 2.4 M NaCl, $5\text{ }\mu\text{g}$ RSV βp4 RT, and $5\text{ }\mu\text{g}$ RSV protease. Incubations were for 1 to 16 hours at 4°C . Digestion products were analyzed by SDS-PAGE and by assay for recovery of RT activity.

Assay of RNase H Activity of RSV RT

The RNase H activity of RSV RT was determined by monitoring the solubilization of [^3H]poly(A) in [^3H]poly(A)•poly(dT). Reaction mixtures ($50\text{ }\mu\text{l}$)

contained 50 mM Tris-HCl (pH 8.4), 20 mM KCl, 10 mM MgCl₂, 10 μM each of [³H] poly(A) (300 cpm/pmole) and poly(dT) in [³H]poly(A)•poly(dT) and 10 mM dithiothreitol. Reaction mixtures were incubated at 37°C for 20 minutes. Incubations were terminated by the addition of 80 μl of 20% (W/V) TCA and 10 μl of 1 mg/ml tRNA, and after centrifugation the amount of [³H]poly(A) solubilized was determined by counting the supernatant in aqueous scintillation fluid. One unit of RNase H activity was the amount of enzyme that solubilized one nmole of [³H]poly(A) in 20 minutes at 37°C.

Assay of DNA Polymerase Activity with Poly(C)•Oligo(dG)₁₂₋₁₈

The RNA-dependent DNA polymerase activity of RSV RT was determined by monitoring the synthesis of acid insoluble [³H]poly(dG) from poly(C)•oligo(dG). Reaction mixtures (50 μl) contained 50 mM Tris-HCl (pH 8.4), 50 mM KCl, 10 mM MgCl₂, 0.5 mM poly(C), 0.2 mM oligo(dG)₁₂₋₁₈, 0.5 mM [³H]dGTP (40 cpm/pmole), and 10 mM dithiothreitol. Reactions were incubated at 37°C for 10 minutes and labeled products were acid precipitated on GF/C glass filters that were counted in a scintillation counter. One unit of DNA polymerase activity was the amount of enzyme that incorporated one nmole of [³H]dGTP in 10 minutes at 37°C.

RESULTS AND DISCUSSION

Alternative Methods of Generating RSV αβ RT

Examples 1 through 6 demonstrated that RNase H⁻ forms of avian RT are more efficient than RNase H⁺ RT in copying mRNA. The studies presented in this Example were designed to determine the efficiency of mRNA copying by RSV

RNase H⁻ $\alpha\beta$ RT that was generated by co-expression of the RSV α and β genes in several expression systems.

Expression in E. coli. Small amounts of soluble and active α , $\beta\beta$, $\beta p4\beta p4$, and $\alpha\beta$ RSV RT have been purified from *E. coli* (Alexander, F., *et al.*, *J. Virol.* 61: 534 (1987); Weis, J.H., and Salstrom, J.S., U.S. Patent No. 4,663,290 (1987); Soltis, D.A. and Skalka, A.M., *Proc. Nat. Acad. Sci. USA* 85:3372 (1988); and Cherhov, A.P., *et al. Biomed Sci.* 2: 49 (1991)). However, most of the RSV RT expressed in *E. coli* in these previous reports was in an insoluble form.

The present efforts to express amounts of RSV RT proteins easily purified from *E. coli* are documented in the Materials and Methods section of this Example. In general, similar results to those previously published were obtained; that is, most of the RSV RT protein expressed in *E. coli* was insoluble, and only small amounts of RT activity were observed. Because of low RT levels, no attempts were made to purify RSV RT expressed in *E. coli*.

Expression of the RSV RT $\beta p4$ or β gene in cultural insect cells. Heterodimeric p66/p51 HIV RT has been produced in *E. coli* and yeast host cells when the cloned gene for HIV p66 was expressed (Lowe, D.M., *et al.*, *Biochemistry* 27:8884 (1988); Muller, B., *et al.*, *J. Biol. Chem.* 264:13975 (1989); and Barr, P.J., *et al.*, *BioTechnology* 5:486 (1987)). Formation of the heterodimer occurs by proteolytic processing of p66/p66 by endogenous host proteases. In contrast, expression of the gene for HIV RT p66 in cultured insect cells yielded only p66/p66 homodimer (Kawa, S., *et al.*, *Prot. Expression and Purification* 4:298 (1993)).

In the present studies, expression of the gene for RSV RT $\beta p4$ in cultured insect cells was similarly found to result exclusively in the production of homodimer $\beta p4\beta p4$ (data not shown); little processing to $\alpha\beta$ or α was observed. When the RSV RT β gene was expressed in these cells, however, all three forms

of RSV RT were produced (Table 6). Most of the RT present in the cells was $\beta\beta$ (50-80%); a small amount of $\alpha\beta$ was produced (~10%); and even α was obtained (10-40%). These results suggest that endogenous host proteases in cultured insect cells proteolyze $\beta\beta$, but not $\beta p4\beta p4$, to $\alpha\beta$. The RSV $\alpha\beta$ RT generated by proteolysis of $\beta\beta$ had much lower functional activity than $\alpha\beta$ generated by co-expression of the RSV RT α and β genes (Table 7).

Table 6. *Expression Levels of RSV RTs in Insect Cells Infected With RSV RT β Gene*

Infection No.	Amount of RT Present per 30 grams of Insect Cells					
	RSV $\beta\beta$ RT		RSV $\alpha\beta$ RT		RSV α RT	
	mg	%	mg	%	mg	%
1	0.7	54	0.1	8	0.5	38
2	0.12	79	0.017	11	0.015	10
3	0.55	83	0.05	8	0.06	9

Expression of RSV RT $\beta p4$ in insect larvae. Baculovirus bearing the RSV RT $\beta p4$ gene was also used to infect live larvae by physical injection of virus. The level of proteases present in larvae and in larval extracts is much higher than in cultured insect cells. Processing of $\beta p4$ to multiple forms of proteolyzed RT was observed, including processing to α . The major species of RT that could be purified from these extracts had four major bands that migrated on SDS-PAGE at 97 kDa (His-tagged β), 87 kDa (proteolyzed β), 67 kDa (partially processed His-tagged α), and 62 kDa (α with His tag removed by proteolysis). This RSV RT had a specific activity of 55,000 units/mg protein, comparable to RSV $\alpha\beta$ RT. In the functional activity assay (Example 3), however, the RT purified from larvae had 85% and 60% of the total product and full length product functional activity, respectively, of RSV $\alpha\beta$ RT generated by co-expression of α and β (Table 7).

Table 7. Specific and Functional Activities of Various Forms of RSV RTs Expressed in Insect Cells

RT Form Isolated												
Gene(s) Expressed	RSV αβ RT			RSV ββ RT			RSV α RT			RSV βp4βp4 RT		
	Specific Activity ^a (U/mg)	Functional Activity ^b		Specific Activity (U/mg)	Functional Activity		Specific Activity (U/mg)	Functional Activity		Spec. Act. (U/mg)	Functional Activity	
		Total (ng/μg) ^c	Full-length (ng/μg) ^d		Total (ng/μg)	Full-length (ng/μg)		Total (ng/μg)	Full-length (ng/μg)			
α and β	53,191	4,092	NP ^e	NP	NP	ND ^f	ND	NP	NP	NP	NP	
β	25,113	1,098	15,819	584	41	ND	ND	NP	NP	NP	NP	
βp4	NP	NP	NP	NP	NP	NP	NP	NP	15,984	450	67	
α	NP	NP	NP	NP	NP	48,759	272	6	NP	NP	NP	

Specific activity = units (U), D.T. activity = (ng/μg), NP = not present

^aSpecific activity = units (U) RT activity/mg RT protein in poly(A)•oligo(dT) assay (Houts *et al.*, *J. Virol.* 29:517 (1979)).

^bFunctional activity established with 7.5 Kb RNA as described in Example 3.

^cMass of total reverse transcribed product, ng of product produced per μg of RT used.

^dMass of full-length reverse transcribed product, ng of product produced per μg of RT used.

^e"NP" = RT form not produced by insect cells expressing the indicated gene.

^f"ND" = RT form produced by insect cells, but not analyzed in present studies.

Co-expression of RSV RT β p4 and RSV protease p15 in cultured insect cells. Heterodimeric p66/p51 HIV RT has been produced efficiently in *E. coli* by co-expression of HIV protease and HIV RT p66 (Mizuuchi, V., *et al.*, *Arch. Biochem Biophys.* 273:347 (1989) and Le Grice, S.F.J. and Gruninger-Leitch, F., *Eur. J. Biochem.* 187:307 (1990)). Co-expression of the viral protease increased the overall efficiency of converting p66/p66 to heterodimer. As described above in the Materials and Methods section of this Example, co-expression of RSV RT β and RSV protease p15 genes in *E. coli* did not result in enhanced production of RSV $\alpha\beta$ RT. Similarly, co-expression of RSV RT β p4 and RSV protease p15 in cultured insect cells did not appreciably enhance the formation of RSV $\alpha\beta$ RT.

Processing of RSV RT β p4 with RSV protease p15 in vitro. Heterodimeric p66/p51 HIV RT has been produced from p66 purified from *E. coli* by treatment with HIV protease *in vitro* (Chattopadhyay, D., *et al.*, *J. Biol. Chem.* 267:14227 (1992)). Purified RSV RT β p4 was treated with RSV protease p15 to generate RSV $\alpha\beta$ RT. This approach was successful in generating some $\alpha\beta$ from β p4 based upon SDS-PAGE analysis, but several difficulties were encountered. First, contrary to what was observed with viral protease treatment of HIV p66 RT, processing of β p4 to $\alpha\beta$ did not always stop at $\alpha\beta$, as α in excess of β was formed as proteolysis proceeded. Second, significant loss of DNA polymerase activity was observed during proteolysis, suggesting RSV RT was partially inactivated by the acid pH reaction conditions required by RSV protease.

Processing of RSV RT β p4 with chymotrypsin in vitro. Heterodimeric p66/p51 HIV RT has also been produced from p66 by limited proteolysis with α -chymotrypsin (Lowe, D.M., *et al.*, *Biochemistry* 27:8884 (1988)). This approach was tried unsuccessfully with RSV RT β p4. We found the digestion of β p4 with α -chymotrypsin was difficult to control, and proteolysis was observed to not stop at $\alpha\beta$, but to proceed to conversion of β p4 to α .

Mixing of RSV RT α and β (in vitro) to generate $\alpha\beta$. Heterodimeric p66/p51 HIV RT has been produced by mixing separate crude cell lysates containing p51 alone and p66 alone (Stahlhut, M., *et al.*, *Protein Expression and Purification* 5:614 (1994)). Mixing of the separate subunits results in formation of a 1:1 molar complex of p66/p51. In contrast, mixing of purified RSV α RT with purified RSV β RT at approximately a 1:1 molar ratio did not result in the formation of an $\alpha\beta$ complex. These results are consistent with the notion that the RSV subunits, once folded separately in an active conformation, prefer to remain separate when mixed.

Relative Ability of Various Forms of RSV RT to Copy RNA

A comparison was made of the ability of four different forms of RSV RT ($\alpha\beta$, $\beta\beta$, α , and $\beta p4\beta p4$) to copy RNA. The RNase H active site of each subunit in these enzymes was mutated to eliminate RNase H activity. Each RT was expressed in cultured insect cells and purified by methods described above and in Example 1. Two RNAs were used for comparison: synthetic homopolymer poly(A) and 7.5-Kb mRNA. With poly(A)•oligo(dT) as template-primer, a specific activity was calculated by determining an initial rate of poly(dT) synthesis catalyzed at limiting enzyme concentration, and then normalizing the rate to a given mass (mg) of RT in the reaction. This specific activity simply represents the ability of a given RT to incorporate a single deoxynucleotide with an artificial template, and does not necessarily represent the ability of the enzyme to copy heteropolymeric RNA. With 7.5-Kb RNA as template, the ability of the RTs to make a full-length copy of a long heteropolymeric RNA was assessed (see Example 3 for details). The results are shown above in Table 7.

Two different forms of $\alpha\beta$ were characterized in Table 7. One form was generated as the result of the expression of the RSV RT β gene and subsequent proteolytic processing in host insect cells, and had reduced specific and functional activity. The other form of $\alpha\beta$ was generated by co-expression of the RSV RT

5 α and β genes. This form of $\alpha\beta$ had a similar specific activity to α , approximately 50,000 units/mg, and had a higher specific activity than either $\beta\beta$ or $\beta p4\beta p4$ (approximately 16,000 units/mg). Comparison of the functional activity of this $\alpha\beta$ to other RT forms showed a much more dramatic contrast. RSV $\alpha\beta$ RT produced 7, 9 and 15 times more total cDNA per mass of enzyme than $\beta\beta$, $\beta p4\beta p4$ and α , respectively, from 7.5-Kb RNA. Even greater differences were observed when yield of full length product was assessed: RSV $\alpha\beta$ RT produced 21, 13 and 146 times more full length product per mass of enzyme than $\beta\beta$, $\beta p4\beta p4$ and α , respectively. RSV $\alpha\beta$ RT produced by co-expression of the RSV RT α H⁺ and β H⁺ genes is therefore much more efficient in copying mRNA than is any other form of RSV RT prepared by analogous methods.

Evidence that Only the α Subunit in RSV $\alpha\beta$ RT is Active

5 Selective DNA polymerase active site mutagenesis of the HIV RT heterodimer p66/p51 has shown that only the DNA polymerase active site of p66 is crucial for DNA polymerase activity (LeGrice, S.F.J., *et al.*, *The EMBO Journal* 10: 3905 (1991) and Hostomsky, Z., *et al.*, *J. Virol.* 66: 3179 (1992)). The p51 subunit in the HIV p66/p51 heterodimer apparently assumes a conformation which does not have a substrate binding cleft and therefore does not participate directly in dNTP binding and incorporation (Jacob-Molina, A., *et al.*, *Proc Natl. Acad Sci USA* 90: 6320 (1993)).

20 In the present studies, the same question was asked for RSV $\alpha\beta$ RT. In this case, since each subunit contains both a DNA polymerase and RNase H active site, combinations of DNA polymerase mutants alone and RNase H mutants alone were characterized. The results are shown in Table 8.

Table 8. DNA Polymerase and RNase H Activities of Various Forms of RSV RT.

RSV RT Form	Specific Activity (Units/mg)		Ratio of DNA Polymerase/RNase H
	DNA Polymerase	RNase H	
$\alpha H^+/\beta H^+$	52,095	924	56.4
$\alpha H^+/\beta H^-$	53,191	<0.1	--
$\alpha H^-/\beta H^-$	48,250	760	64.5
$\alpha H^-/\beta H^+$	50,909	4.5	11,313
$\beta H^+/\beta H^-$	15,819	<0.1	--
αH^-	48,759	<0.1	--
$\alpha pol^+/\beta pol^+$	52,095	924	56.4
$\alpha pol^-/\beta pol^-$	50	600	0.08
$\alpha pol^+/\beta pol^-$	57,500	900	63.9
$\alpha pol^-/\beta pol^+$	1,143	629	1.82

As shown in Table 8, when the RNase H active site in both subunits was mutated to eliminate RNase H activity, RNase H activity in purified enzyme was reduced to below detectable levels, while DNA polymerase activity was unchanged. When the RNase H active site in β was altered with the same mutation and the α RNase H active site was wild type, the enzyme was similar to wild type in polymerase and RNase H activity. In contrast, mutation of the α subunit in RNase H, but not the β subunit, resulted in a 200-fold reduction in RNase H activity. Therefore, in the case of RSV $\alpha\beta$ RT, the RNase H domain of the large subunit (β) folds in an inactive conformation. Examination of DNA polymerase active site mutants in RSV $\alpha\beta$ RT revealed the same results (Table 8). The α subunit, but not the β subunit, supplies the DNA polymerase catalytic activity. So, in contrast to HIV p66/p51 RT, the smaller RSV $\alpha\beta$ RT subunit, not the larger, maintains enzymatic activity. The common element in both enzymes is that the subunit possessing DNA polymerase and RNase H domains folds in an active

conformation, while subunits missing the RNase H domain or possessing an additional domain (integrase), fold in an inactive conformation.

In sum, these results demonstrate that only the $\alpha\beta$ form of ASLV RT copies mRNA efficiently. In addition, the present results indicate that expression of ASLV RT in *E. coli* results in little or no production of active RT, while expression and activity increase dramatically by cloning and expression of ASLV RT in insect cells and yeast. Finally, by placing point mutations in either the DNA polymerase or the RNase H active site of the α or β subunit of RSV $\alpha\beta$ RT, it has now been discovered that the DNA polymerase and RNase H catalytic activities of the RSV RT $\alpha\beta$ heterodimer reside in the α subunit alone.

Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious to one of ordinary skill in the art that the same can be performed by modifying or changing the invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Gerard, Gary F.
Smith, Michael
Chatterjee, Deb

(ii) TITLE OF INVENTION: Compositions and Methods for Reverse
Transcription of Nucleic Acid Molecules

(iii) NUMBER OF SEQUENCES: 23

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.
(B) STREET: 1100 New York Avenue, Suite 600
(C) CITY: Washington
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 20005

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: (To be assigned)
(B) FILING DATE: (Herewith)
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/049,874
(B) FILING DATE: 17-JUN-1997
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/044,589
(B) FILING DATE: 22-APR-1997

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Esmond, Robert W.
(B) REGISTRATION NUMBER: 32,893
(C) REFERENCE/DOCKET NUMBER: 0942.4330002

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-371-2600
(B) TELEFAX: 202-371-2540

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGGAGACCCA TATGACTGTT GCGCTACATC TGGCT

35

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AACGCGACAG GTTAACAGCG CGCAAATCAT GCAG

34

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CACACACAGG TACCCTCTCG AAAAGTTAAA CC

32

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CACACACACT CGAGTTATGC AAAAAGAGGG CTCGCCTCAT C

41

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGACCCACTG TCTTTACCGC GGCCTCCTCA AGCACC

36

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CACACACACC CGGGTTAATA CGCTTGGAAG GTGGC

35

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

5 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CACACACATC ATGACTGTTG CGCTACATCT G

31

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CACACACAGG TACCCTCTCG AAAAGTTAA

29

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CACACACAGA GGAATTCAGT GATGGTGATG GTGATGTGCA AAAAGAGG

48

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACTGGAATTC ATGCCAATCC ATCACCATCA CCATCACCCG T

41

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15 ACGTGTGCGAC CATATGGATG ACTAGGTGAA ACGGGTGATG G

41

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

20 ACTGGAATTC ATGCCAATCC ATCACCATCA CCATCACCCG TTTCACCTAG TCATCCATAT

60

25 GGTCGACACG T

71

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

5

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GACTAGTTCT AGATCGCGAG CGGCCGCCCA TTA ACTCTCG TTGGCAGC

48

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

10

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TCGACCCACG CGTCCG

16

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

20

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGGACGCGTG GG

12

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

5 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AUGGAGAUCU CUGAATTCAT GACTGTTGCG CTACATCTGG CT

42

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATTATTCATA TGACTGTTGC GCTACATCTG GC

32

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TACGATCTCT CTCCAGGCCA TTTT

24

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ACTCGAGCAG CCCGGAACC TTG

24

(2) INFORMATION FOR SEQ ID NO:20:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 48 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

15 ATTACCCGGG AGGATATCAT ATGTTAGCGA TGACAATGGA ACATAAAG

48

(2) INFORMATION FOR SEQ ID NO:21:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 36 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ATATGTCGAC TCACAGTGGC CCTCCCTATA AATTTG

36

25 (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid

(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

5 TATTAGGATC CCATGACTGT TCGCTACAT CTGGC

35

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCAATCCTTG AGCTCTAAGA CCATCAGGG

29

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156

WHAT IS CLAIMED IS:

1. A composition for use in reverse transcription of a nucleic acid molecule, said composition comprising two or more polypeptides having reverse transcriptase activity.

5 2. The composition of claim 1, wherein said polypeptides are obtained from different sources.

3. The composition of claim 1, wherein the transcription pause site of each of said polypeptides is different from that of each of the other polypeptides in said composition.

10 4. The composition of claim 1, wherein said polypeptides are reduced or substantially reduced in RNase H activity.

15 5. The composition of claim 4, wherein said polypeptides are selected from the group consisting of M-MLV H⁻ reverse transcriptase, RSV H⁻ reverse transcriptase, AMV H⁻ reverse transcriptase, RAV H⁻ reverse transcriptase, MAV H⁻ reverse transcriptase and HIV H⁻ reverse transcriptase, and derivatives, variants, fragments or mutants thereof.

20 6. The composition of claim 5, wherein said AMV H⁻ reverse transcriptase is selected from the group consisting of AMV α H⁻/ β H⁻ reverse transcriptase, AMV α H⁻/ β H⁺ reverse transcriptase, AMV β H⁻/ β H⁻ reverse transcriptase, AMV β H⁺/ β H⁻ reverse transcriptase, AMV β p4/ β p4 reverse transcriptase, and AMV α H⁻ reverse transcriptase, and derivatives, variants, fragments or mutants thereof.

7. The composition of claim 5, wherein said RSV H⁻ reverse transcriptase is selected from the group consisting of RSV αH⁻/βH⁻ reverse transcriptase, RSV αH⁻/βH⁺ reverse transcriptase, RSV βH⁻/βH⁻ reverse transcriptase, RSV βH⁺/βH⁻ reverse transcriptase, RSV βp4/βp4 reverse transcriptase, and RSV αH⁻ reverse transcriptase, and derivatives, variants, fragments or mutants thereof.

8. The composition of claim 1, wherein said polypeptides are selected from the group consisting of *Taq*, *Tne*, *Tma*, *Pfu*, VENT™, DEEPVENT™ and *Tth* DNA polymerases, and mutants, fragments, variants and derivatives thereof.

9. The composition of claim 1, wherein said polypeptides are present in said composition at working concentrations.

10. A method for reverse transcription of one or more nucleic acid molecules comprising

(a) mixing one or more nucleic acid templates with two or more polypeptides having reverse transcriptase activity; and

(b) incubating said mixture under conditions sufficient to make one or more first nucleic acid molecules complementary to all or a portion of said one or more templates.

11. The method of claim 10, wherein said nucleic acid template is a messenger RNA molecule or a population of mRNA molecules.

12. The method of claim 10, said method further comprising incubating said one or more first nucleic acid molecules under conditions sufficient to make one or more second nucleic acid molecules complementary to all or a portion of said one or more first nucleic acid molecules.

13. A cDNA molecule made according to the method of claim 10.

14. A cDNA molecule made according to the method of claim 12.

15. A method for amplifying one or more nucleic acid molecules, said method comprising

5 (a) mixing one or more nucleic acid templates with two or more polypeptides having reverse transcriptase activity and one or more DNA polymerases; and

(b) incubating said mixture under conditions sufficient to amplify one or more nucleic acid molecules complementary to all or a portion of said one or more templates.

16. A method for amplifying one or more nucleic acid molecules, said method comprising

15 (a) mixing one or more nucleic acid templates with two or more polypeptides having reverse transcriptase activity and DNA polymerase activity; and

(b) incubating said mixture under conditions sufficient to amplify one or more nucleic acid molecules complementary to all or a portion of said one or more nucleic acid templates.

20 17. A nucleic acid molecule amplified according to the method of claim 15 or claim 16.

18. A vector comprising the cDNA molecule of claim 14.

19. The vector of claim 18, wherein said vector is an expression vector.

20. A host cell comprising the cDNA molecule of claim 14.

21. A method for sequencing one or more nucleic acid molecules, said method comprising

(a) mixing one or more nucleic acid molecules to be sequenced with one or more primers, two or more polypeptides having reverse transcriptase activity, one or more nucleotides and one or more terminating agents;

(b) incubating said mixture under conditions sufficient to synthesize a population of molecules complementary to all or a portion of said one or more molecules to be sequenced; and

(c) separating said population to determine the nucleotide sequence of all or a portion of said one or more molecules to be sequenced.

22. A kit for use in reverse transcription, amplification or sequencing of a nucleic acid molecule, said kit comprising two or more polypeptides having reverse transcriptase activity.

23. The kit of claim 22, said kit further comprising one or more components selected from the group consisting of one or more nucleotides, one or more DNA polymerases, a suitable buffer, one or more primers and one or more terminating agents.

24. The kit of claim 23, wherein said terminating agent is a dideoxynucleotide.

25. The kit of claim 23, wherein two or more of the components of said kit are present as a mixture or are present as separate components.

26. A method of producing an ASLV reverse transcriptase, said method comprising

(a) obtaining a host cell comprising one or more nucleic acid sequences encoding one or more subunits of ASLV reverse transcriptase; and

(b) culturing said host cell under conditions sufficient to produce said ASLV reverse transcriptase subunits.

27. The method of claim 26, wherein said one or more nucleic acid sequences encoding one or more subunits of ASLV reverse transcriptase are contained in one or more vectors.

28. The method of claim 26, wherein said ASLV reverse transcriptase subunits are selected from the group consisting of one or more α subunits, one or more β subunits, and one or more β p4 subunits, of one or more ASLV reverse transcriptases, and derivatives, variants, fragments or mutants thereof.

29. The method of claim 26, wherein said subunits are one or more α subunits.

30. The method of claim 26, wherein said subunits are one or more β subunits.

31. The method of claim 26, wherein said subunits are one or more β p4 subunits.

32. The method of claim 26, wherein said subunits are one α subunit and one β subunit of one or more ASLV reverse transcriptases.

33. The method of claim 26, wherein said subunits are co-expressed to form an ASLV reverse transcriptase and wherein said ASLV reverse transcriptase is isolated from said host cell.

5 34. The method of claim 26, wherein said subunits of ASLV reverse transcriptase are isolated and then mixed to form an ASLV reverse transcriptase.

35. The method of claim 30, wherein said β subunits form an ASLV reverse transcriptase comprising two β subunits.

36. The method of claim 32, wherein said α and β subunits form an ASLV reverse transcriptase comprising an α and a β subunit.

10 37. The method of claim 26, wherein one or more of said subunits have been modified to reduce or substantially reduce the RNase H activity in said subunits.

38. The method of claim 26, wherein said subunits are encoded by one or more nucleotide sequences contained on the same or on different vectors.

15 39. The method of claim 26, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

40. The method of claim 26, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

20 41. An ASLV reverse transcriptase produced according to the method of claim 26.

42. The ASLV reverse transcriptase of claim 41, wherein said ASLV reverse transcriptase is selected from the group consisting of an ASLV $\alpha\beta$ reverse transcriptase, an ASLV $\beta\beta$ reverse transcriptase, an ASLV $\beta p4\beta p4$ reverse transcriptase, and an ASLV α reverse transcriptase.

5 43. The ASLV reverse transcriptase of claim 41, wherein said ASLV reverse transcriptase is reduced or substantially reduced in RNase H activity.

44. The ASLV reverse transcriptase of claim 41, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

10 45. The ASLV reverse transcriptase of claim 41, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

46. An isolated nucleic acid molecule comprising a nucleotide sequence encoding one or more subunits of ASLV reverse transcriptase.

15 47. The nucleic acid molecule of claim 46, wherein said molecule encodes one or more α subunits of ASLV reverse transcriptase, or a derivative, fragment or mutant thereof.

48. The nucleic acid molecule of claim 46, wherein said molecule encodes one or more β subunits of ASLV reverse transcriptase, or a derivative, fragment or mutant thereof.

20 49. The nucleic acid molecule of claim 46, wherein said molecule encodes one or more $\beta p4$ subunits of ASLV reverse transcriptase, or a derivative, fragment or mutant thereof.

50. The nucleic acid molecule of claim 46, wherein said molecule encodes α and β ASLV reverse transcriptase subunits, or derivatives, variants, fragments or mutants thereof.

51. A vector comprising the nucleic acid molecule of claim 46.

52. A host cell comprising the nucleic acid molecule of claim 46.

53. The vector of claim 51, wherein said vector is plasmid pDABH-His.

54. The host cell of claim 52, wherein said host cell is *E. coli* DH10B(pDABH-His).

55. The isolated nucleic acid molecule of claim 46, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

56. The isolated nucleic acid molecule of claim 46, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

57. A method for producing one or more cDNA molecules by reverse transcription of one or more nucleic acid templates comprising

(a) mixing one or more nucleic acid templates with a ASLV RT comprising one or more subunits; and

(b) incubating said mixture under conditions sufficient to make one or more first nucleic acid molecules complementary to all or a portion of said one or more templates.

58. The method of claim 57, wherein said subunits are one or more α subunits, one or more β subunits, one or more $\beta p4$ subunits, or a combination thereof.

5 59. The method of claim 57, wherein said ASLV reverse transcriptase is selected from the group consisting of an ASLV $\alpha\beta$ reverse transcriptase, an ASLV $\beta\beta$ reverse transcriptase, an ASLV $\beta p4\beta p4$ reverse transcriptase, and an ASLV α reverse transcriptase.

60. The method of claim 57, wherein said nucleic acid template is a mRNA molecule or a population of mRNA molecules.

10 61. The method of claim 57, wherein said method further comprises incubating said one or more first nucleic acid molecules under conditions sufficient to make one or more second nucleic acid molecules complementary to all or a portion of said one or more first nucleic acid molecules.

15 62. The method of claim 61, wherein said first and said second nucleic acid molecules form a double stranded DNA molecule.

63. The method of claim 62, wherein said double stranded DNA molecule is a full-length cDNA molecule.

64. A cDNA molecule made according to the method of claim 57.

65. A cDNA molecule made according to the method of claim 61.

20 66. A vector comprising the cDNA molecule of claim 65.

67. The vector of claim 66, wherein said vector is an expression vector.

68. A host cell comprising the cDNA molecule of claim 65.

69. The method of claim 61, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

5 70. The method of claim 61, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

71. A method for amplifying one or more nucleic acid molecules comprising

(a) mixing one or more nucleic acid templates with one or more ASLV RTs comprising one or more subunits and optionally with one or more DNA polymerases; and

(b) incubating said mixture under conditions sufficient to amplify one or more nucleic acid molecules complementary to all or a portion of said one or more templates.

15 72. The method of claim 71, wherein said subunits are one or more α subunits, one or more β subunits, one or more β p4 subunits, or a combination thereof.

20 73. The method of claim 71, wherein said ASLV reverse transcriptase is selected from the group consisting of an ASLV $\alpha\beta$ reverse transcriptase, an ASLV $\beta\beta$ reverse transcriptase, an ASLV β p4 β p4 reverse transcriptase, and an ASLV α reverse transcriptase.

74. The method of claim 71, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

75. The method of claim 71, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

5 76. A method for sequencing one or more nucleic acid molecules comprising

(a) mixing one or more nucleic acid molecules to be sequenced with one or more primers, an ASLV RT comprising one or more subunits, one or more nucleotides and one or more terminating agents;

10 (b) incubating said mixture under conditions sufficient to synthesize a population of nucleic acid molecules complementary to all or a portion of said one or more nucleic acid molecules to be sequenced; and

15 (c) separating said population of nucleic acid molecules to determine the nucleotide sequence of all or a portion of said one or more nucleic acid molecules- to be sequenced.

77. The method of claim 76, wherein said subunits are one or more α subunits, one or more β subunits, one or more β p4 subunits, or a combination thereof.

20 78. The method of claim 76, wherein said ASLV reverse transcriptase is selected from the group consisting of an ASLV $\alpha\beta$ reverse transcriptase, an ASLV $\beta\beta$ reverse transcriptase, an ASLV β p4 β p4 reverse transcriptase, and an ASLV α reverse transcriptase.

79. The method of claim 76, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

80. The method of claim 76, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

81. A kit comprising one or more ASLV RT subunits, or one or more derivatives, variants, fragments or mutants thereof.

82. The kit of claim 81, wherein said ASLV RT subunits are one or more ASLV RT α subunits, one or more ASLV RT β subunits, one or more ASLV RT β p4 subunits, or a combination thereof.

83. The kit of claim 81, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

84. The kit of claim 81, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

85. A kit comprising an ASLV RT, wherein said ASLV RT is selected from the group consisting of an ASLV $\alpha\beta$ RT, an ASLV $\beta\beta$ RT, an ASLV β p4 β p4 RT, and an ASLV α RT, or a derivative, fragment or mutant thereof.

86. The kit of claim 85, wherein said ASLV RT comprises an α and a β subunit.

87. The kit of claim 85, wherein said ASLV reverse transcriptase is an RSV reverse transcriptase.

88. The kit of claim 85, wherein said ASLV reverse transcriptase is an AMV reverse transcriptase.

89. A method for producing one or more cDNA molecules by reverse transcription of one or more nucleic acid templates comprising

(a) mixing one or more nucleic acid templates with one or more polypeptides having reverse transcriptase activity; and

(b) incubating said mixture at a temperature of about 50°C or greater and under conditions sufficient to make one or more first nucleic acid molecules complementary to all or a portion of said one or more templates.

90. The method of claim 89, wherein said temperature is 60°C or greater.

91. The method of claim 89, wherein said temperature ranges from about 50°C to about 70°C.

92. The method of claim 89, wherein said temperature ranges from about 55°C to about 65°C.

93. The method of claim 89, wherein said nucleic acid template is an RNA or DNA molecule.

94. The method of claim 93, wherein said RNA molecule is an mRNA molecule or a polyA⁺ RNA molecule.

95. The method of claim 89, wherein said nucleic acid template is a population of mRNA molecules.

96. The method of claim 94, wherein said first nucleic acid molecule is a full length cDNA molecule.

97. The method of claim 89, further comprising incubating said one or more first nucleic acid molecules under conditions sufficient to make one or more second nucleic acid molecules complementary to all or portion of said one or more first nucleic acid molecules.

5 98. The method of claim 97, wherein said first and said second nucleic acid molecules are DNA molecules.

99. The method of claim 98, wherein said first and said second DNA molecules form a double stranded DNA molecule.

100. The method of claim 99, wherein said double stranded DNA molecule is a full length cDNA molecule.

101. The method of claim 89, wherein said one or more polypeptides having reverse transcriptase activity are reduced or substantially reduced in RNase H activity.

102. The method of claim 89, wherein said one or more polypeptides having reverse transcriptase activity is one or more ASLV reverse transcriptases comprising one or more subunits.

103. The method of claim 102, wherein said one or more ASLV reverse transcriptases is one or more RSV reverse transcriptases.

104. The method of claim 102, wherein said one or more ASLV reverse transcriptases is one or more AMV reverse transcriptases.

105. The method of claim 102, wherein said ASLV RT subunits are one or more ASLV RT α subunits, one or more ASLV RT β subunits, one or more ASLV RT β p4 subunits, or a combination thereof.

5 106. The method of claim 102, wherein said ASLV RT is selected from the group consisting of an ASLV $\alpha\beta$ RT, an ASLV $\beta\beta$ RT, an ASLV β p4 β p4 RT, and an ASLV α RT, and derivatives, variants, fragments or mutants thereof.

107. The method of claim 102, wherein said ASLV RT is ASLV $\alpha\beta$ RT, or a derivative, variant, fragment or mutant thereof.

10 108. The method of claim 102, wherein said one or more subunits are reduced or substantially reduced in RNase H activity.

109. A nucleic acid molecule produced according to the method of claim 89.

110. A nucleic acid molecule produced according to the method of claim 97.

15 111. The nucleic acid molecule of claim 110, wherein said nucleic acid molecule is a full-length cDNA molecule.

112. A vector comprising the nucleic acid molecule of claim 110.

113. The vector of claim 112, wherein said vector is an expression vector.

20 114. A host cell comprising the nucleic acid molecule of claim 109.

115. A host cell comprising the nucleic acid molecule of claim 110.

116. A host cell comprising the nucleic acid molecule of claim 110.

Compositions and Methods for Reverse Transcription of Nucleic Acid Molecules

ABSTRACT

The present invention is generally related to compositions and methods for the reverse transcription of nucleic acid molecules, especially messenger RNA molecules. Specifically, the invention relates to compositions comprising mixtures of polypeptides having reverse transcriptase (RT) activity, and to methods of producing, amplifying or sequencing nucleic acid molecules (particularly cDNA molecules) using these compositions or polypeptides, particularly at temperatures above about 55°C. The invention also relates to nucleic acid molecules produced by these methods, to vectors and host cells comprising these nucleic acid molecules, and to the use of such nucleic acid molecules to produce desired polypeptides. The invention also relates to methods for producing Rous Sarcoma Virus (RSV) and Avian Myeloblastosis Virus (AMV) RTs or other Avian Sarcoma-Leukosis Virus (ASLV) RTs (α and/or β subunits thereof), to isolated nucleic acid molecules encoding such RSV RT, AMV RT or other ASLV RT subunits, to vectors and host cells comprising these isolated nucleic acid molecules and to RSV RT, AMV RT and other ASLV RT subunits produced by these methods. The invention further relates to nucleic acid molecules encoding recombinant heterodimeric RT holoenzymes, particularly heterodimeric RSV RTs, AMV RTs or other ASLV RTs (which may be $\alpha\beta$ RTs, $\beta\beta$ RTs, or α RTs), vectors (particularly baculovirus vectors) and host cells (particularly insect and yeast cells) comprising these nucleic acid molecules, methods for producing these heterodimeric RTs and heterodimeric RTs produced by these methods. The invention also relates to kits comprising the compositions, polypeptides, or RSV RTs, AMV RTs or other ASLV RTs of the invention.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

GERARD et al.

Appl. No. (To be assigned; Divisional of
U.S. Appl. No. 09/064,057)

Filed: (Herewith)

For: **Compositions and Methods for
Reverse Transcription of Nucleic
Acid Molecules**

Art Unit: (To be assigned)

Examiner: (To be assigned)

Atty. Docket: 0942.4330003/RWE/BJD

Request to Open New Disk File

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Applicants request that a new disk file be opened for the above-cited application. The Sequence Listing disk submitted on April 22, 1998, in the parent, U.S. Application No. 09/064,057, filed April 22, 1998, contains the identical sequence information as that in the present application.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Brian J. Del Buono
Agent for Applicants
Registration No. 42,473

Date: February 5, 1999

1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005-3934
(202) 371-2600

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.





Figure 6.






Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21

FIGURE 22




FIGURE 23.




FIGURE 24.




FIGURE 25.




FIGURE 26.




FIGURE 27




FIGURE 2B.




FIGURE 29.




FIGURE 3D.




FIGURE 31.




FIGURE 32.




FIGURE 33.




FIGURE 34.




FIGURE 35.




FIGURE 36




FIGURE 37.




FIGURE 3B:





Fig 39



Parameter	Estimate	Standard Error	t-Statistic	p-Value	95% Confidence Interval
Intercept	1.2345	0.0123	100.34	0.0000	1.2100 - 1.2590
Variable 1	0.4567	0.0234	19.50	0.0000	0.4099 - 0.5035
Variable 2	-0.1234	0.0156	-7.91	0.0000	-0.1546 - -0.0922
Variable 3	0.7890	0.0345	22.87	0.0000	0.7199 - 0.8581
Variable 4	0.2345	0.0189	12.40	0.0000	0.2067 - 0.2623
Variable 5	-0.3456	0.0212	-16.30	0.0000	-0.3878 - -0.3034
Variable 6	0.5678	0.0267	21.27	0.0000	0.5144 - 0.6212
Variable 7	0.1234	0.0145	8.44	0.0000	0.0944 - 0.1524
Variable 8	-0.4567	0.0198	-23.07	0.0000	-0.4963 - -0.4171
Variable 9	0.6789	0.0312	21.76	0.0000	0.6165 - 0.7413
Variable 10	0.3456	0.0223	15.49	0.0000	0.3011 - 0.3901
Variable 11	-0.2345	0.0178	-13.18	0.0000	-0.2701 - -0.1989
Variable 12	0.7890	0.0356	22.16	0.0000	0.7178 - 0.8602
Variable 13	0.4567	0.0245	18.64	0.0000	0.4077 - 0.5057
Variable 14	-0.1234	0.0167	-7.39	0.0000	-0.1587 - -0.0881
Variable 15	0.5678	0.0278	20.42	0.0000	0.5122 - 0.6234
Variable 16	0.2345	0.0199	11.78	0.0000	0.1947 - 0.2743
Variable 17	-0.3456	0.0221	-15.64	0.0000	-0.3897 - -0.3015
Variable 18	0.6789	0.0323	21.02	0.0000	0.6143 - 0.7435
Variable 19	0.3456	0.0234	14.77	0.0000	0.3007 - 0.3905
Variable 20	-0.2345	0.0189	-12.40	0.0000	-0.2701 - -0.1989
Variable 21	0.7890	0.0367	21.49	0.0000	0.7156 - 0.8624
Variable 22	0.4567	0.0256	17.84	0.0000	0.4055 - 0.5079
Variable 23	-0.1234	0.0178	-6.93	0.0000	-0.1597 - -0.0871
Variable 24	0.5678	0.0289	19.30	0.0000	0.5100 - 0.6256
Variable 25	0.2345	0.0209	11.22	0.0000	0.1926 - 0.2764
Variable 26	-0.3456	0.0231	-14.96	0.0000	-0.3887 - -0.3025
Variable 27	0.6789	0.0334	20.33	0.0000	0.6129 - 0.7449
Variable 28	0.3456	0.0245	14.11	0.0000	0.2991 - 0.3961
Variable 29	-0.2345	0.0199	-11.78	0.0000	-0.2701 - -0.1989
Variable 30	0.7890	0.0378	20.87	0.0000	0.7139 - 0.8641
Variable 31	0.4567	0.0267	17.11	0.0000	0.4044 - 0.5090
Variable 32	-0.1234	0.0189	-6.52	0.0000	-0.1607 - -0.0861
Variable 33	0.5678	0.0301	18.86	0.0000	0.5077 - 0.6279
Variable 34	0.2345	0.0219	10.71	0.0000	0.1906 - 0.2784
Variable 35	-0.3456	0.0241	-14.34	0.0000	-0.3877 - -0.3035
Variable 36	0.6789	0.0345	19.68	0.0000	0.6124 - 0.7454
Variable 37	0.3456	0.0256	13.46	0.0000	0.2976 - 0.3936
Variable 38	-0.2345	0.0209	-11.22	0.0000	-0.2701 - -0.1989
Variable 39	0.7890	0.0389	20.28	0.0000	0.7101 - 0.8679
Variable 40	0.4567	0.0278	16.42	0.0000	0.4033 - 0.5101
Variable 41	-0.1234	0.0199	-6.20	0.0000	-0.1599 - -0.0869
Variable 42	0.5678	0.0312	18.19	0.0000	0.5066 - 0.6290
Variable 43	0.2345	0.0229	10.24	0.0000	0.1886 - 0.2804
Variable 44	-0.3456	0.0251	-13.77	0.0000	-0.3867 - -0.3045
Variable 45	0.6789	0.0356	19.07	0.0000	0.6104 - 0.7474
Variable 46	0.3456	0.0267	12.94	0.0000	0.2966 - 0.3946
Variable 47	-0.2345	0.0219	-10.71	0.0000	-0.2701 - -0.1989
Variable					




Figure 41




Figure 42




Figure 43




Figure 44




Figure 46




Figure 4b




Figure 47




Figure 48




Figure 49




Figure 50




figure 51




Figure 52




Figure 53





Figure 54



Declaration for Patent Application

Docket Number: 0942.4330002/RWE/BJD

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled Compositions and Methods for Reverse Transcription of Nucleic Acid Molecules, the specification of which is attached hereto unless the following box is checked:

- ☒ was filed on April 22, 1998;
as United States Application Number or PCT International Application Number 09/064,057; and
was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Claimed

_____ (Application No.)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Application No.)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

<u>60/044,589</u> (Application No.)	<u>April 22, 1997</u> (Filing Date)
<u>60/049,874</u> (Application No.)	<u>June 17, 1997</u> (Filing Date)

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56 that became available between the filing date of the prior application and the national or PCT international filing date of this application.

_____ (Application No.)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)
----------------------------	------------------------	--




Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.
1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor		Gary F. GERARD
Inventor's signature		Date 6/2/98
Residence	Frederick, MD 21704	
Citizenship	USA	
Post Office Address 8393 William Drive Frederick, MD 21704		
Full name of second inventor Michael D. SMITH		
Second Inventor's signature		Date 6/2/98
Residence	Rockville, MD 20853	
Citizenship	USA	
Post Office Address 14005 Shippers Lane Rockville, MD 20853		
Full name of third inventor Deb K. CHATTERJEE		
Third inventor's signature		Date 6/2/98
Residence	North Potomac, MD 20878.	
Citizenship	USA	
Post Office Address 6 Forest Ridge Court North Potomac, MD 20878		